

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTALDB1623

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	JUL 02	LMEDLINE coverage updated
NEWS	3	JUL 02	SCISEARCH enhanced with complete author names
NEWS	4	JUL 02	CHEMCATS accession numbers revised
NEWS	5	JUL 02	CA/CAPLUS enhanced with utility model patents from China
NEWS	6	JUL 16	CAPLUS enhanced with French and German abstracts
NEWS	7	JUL 18	CA/CAPLUS patent coverage enhanced
NEWS	8	JUL 26	USPATFULL/USPAT2 enhanced with IPC reclassification
NEWS	9	JUL 30	USGENE now available on STN
NEWS	10	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	11	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	12	AUG 13	CA/CAPLUS enhanced with additional kind codes for granted patents
NEWS	13	AUG 20	CA/CAPLUS enhanced with CAS indexing in pre-1907 records
NEWS	14	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS	15	AUG 27	USPATOLD now available on STN
NEWS	16	AUG 28	CAS REGISTRY enhanced with additional experimental spectral property data
NEWS	17	SEP 07	STN AnaVist, Version 2.0, now available with Derwent World Patents Index
NEWS	18	SEP 13	FORIS renamed to SOFIS
NEWS	19	SEP 13	INPADOCDB enhanced with monthly SDI frequency
NEWS	20	SEP 17	CA/CAPLUS enhanced with printed CA page images from 1967-1998
NEWS	21	SEP 17	CAPLUS coverage extended to include traditional medicine patents
NEWS	22	SEP 24	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	23	OCT 02	CA/CAPLUS enhanced with pre-1907 records from Chemisches Zentralblatt
NEWS	24	OCT 19	BEILSTEIN updated with new compounds
NEWS	25	NOV 15	Derwent Indian patent publication number format enhanced
NEWS	26	NOV 19	WPIX enhanced with XML display format

NEWS EXPRESS 19 SEPTEMBER 2007: CURRENT WINDOWS VERSION IS V8.2,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 SEPTEMBER 2007.

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS LOGIN Welcome Banner and News Items

NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:58:30 ON 26 NOV 2007

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 15:58:43 ON 26 NOV 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 26 Nov 2007 VOL 147 ISS 23

FILE LAST UPDATED: 25 Nov 2007 (20071125/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> e myllymaki/au

E1	1	MYLLYMAEKI SARI/AU
E2	19	MYLLYMAEKI VESA/AU
E3	0 -->	MYLLYMAKI/AU
E4	1	MYLLYMAKI A/AU
E5	2	MYLLYMAKI ARVO/AU
E6	1	MYLLYMAKI HANNU/AU
E7	1	MYLLYMAKI HILKKA/AU
E8	1	MYLLYMAKI J/AU
E9	2	MYLLYMAKI JUHANI/AU
E10	3	MYLLYMAKI JUKKA/AU
E11	1	MYLLYMAKI JUSSI PETRI/AU
E12	1	MYLLYMAKI LINDA/AU

=> s e2

L1 19 "MYLLYMAEKI VESA"/AU

=> e aksela/au

E1 1 AKSEL VOD L S/AU
E2 2 AKSEL YU Z/AU
E3 0 --> AKSELA/AU
E4 186 AKSELA H/AU
E5 28 AKSELA HELENA/AU
E6 5 AKSELA MAIJA/AU
E7 6 AKSELA R/AU
E8 1 AKSELA RAIMO/AU
E9 1 AKSELA RALMO/AU
E10 1 AKSELA REIGO/AU
E11 46 AKSELA REIJO/AU
E12 201 AKSELA S/AU

=> s e10 or e11

1 "AKSELA REIGO"/AU
46 "AKSELA REIJO"/AU
L2 47 "AKSELA REIGO"/AU OR "AKSELA REIJO"/AU

=> dup remove l1 l2

PROCESSING COMPLETED FOR L1

PROCESSING COMPLETED FOR L2

L3 60 DUP REMOVE L1 L2 (6 DUPLICATES REMOVED)

=> s l3 and ionic

L4 19 S L3
L5 41 S L3
284921 IONIC
511 IONICS
285185 IONIC
(IONIC OR IONICS)
L6 6 (L4 OR L5) AND IONIC

=> d l6 1-6 ibib abs kwic

L6 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:54651 CAPLUS

DOCUMENT NUMBER: 146:144557

TITLE: Production of starch ethers in ionic liquids
in the absence of water

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 20pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2007006848	A1	20070118	WO 2006-FI248	20060712
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,			

GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

PRIORITY APPLN. INFO.:

FI 2005-752

A 20050714

OTHER SOURCE(S): MARPAT 146:144557

AB Starch ethers are prepared by mixing starch with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch with an etherifying agent in the presence of a base to form a starch ether, and subsequently separating the starch ether from the solution, wherein both the dissoln. and the etherification are carried out in the substantial absence of water.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Production of starch ethers in ionic liquids in the absence of water

IN Myllymaeki, Vesa; Aksela, Reijo

AB Starch ethers are prepared by mixing starch with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch with an etherifying agent in the presence of. . .

ST starch dissolving ionic liq etherification

IT Etherification

Ionic liquids

(production of starch ethers in ionic liqs. in absence of water)

IT 79917-90-1, [BMIM]Cl

RL: NUU (Other use, unclassified); USES (Uses)

(ionic liquid; production of starch ethers in ionic liqs. in absence of water)

IT 9005-25-8DP, Starch, ethers

RL: IMF (Industrial manufacture); PREP (Preparation)

(production of starch ethers in ionic liqs. in absence of water)

IT 9057-06-1P, Carboxymethyl starch

RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)

(production of starch ethers in ionic liqs. in absence of water)

L6 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:977559 CAPLUS

DOCUMENT NUMBER: 145:337750

TITLE: Water-insoluble polysaccharide-based composite materials for use in paper and board manufacturing

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo; Sundquist, Anna; Karvinen, Saira Marjatta

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 70pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

-----	-----	-----	-----	-----
WO 2006097571	A1	20060921	WO 2006-FI88	20060315
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
FI 2005000293	A	20060919	FI 2005-293	20050318
CA 2599423	A1	20060921	CA 2006-2599423	20060315
PRIORITY APPLN. INFO.:			FI 2005-293	A 20050318
			WO 2006-FI88	W 20060315

OTHER SOURCE(S): MARPAT 145:337750

AB The invention relates to a composite material based on water-insol. polysaccharide, such as cellulose and chitin. The composite material comprises particles of at least one light scattering material, the surface of which is essentially covered with at least one water-insol. polysaccharide. The invention also relates to a method for preparation of the composite material, and to a paper and board manufacturing process in which the composite material is used as a filler or pigment. Both highly organic products with exceptional heat capacities, as well as cheap high filler products can be manufactured. The composite material significantly improves retention of light scattering fillers in the manufacture of paper and board even without the use of sep. retention aids.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

IN Myllymaeki, Vesa; Aksela, Reijo; Sundquist, Anna; Karvinen, Salla Marjatta

IT Extrusion of plastics and rubbers
Fillers

Ionic liquids

Microparticles

Microspheres

Paper

Paperboard

Pigments, nonbiological

Plastic films

(water-insol. polysaccharide-based composite materials for use in paper and board manufacturing)

L6 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:638995 CAPLUS

DOCUMENT NUMBER: 143:135160

TITLE: Starch depolymerization in ionic liquid solvents

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 21 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005066374	A1	20050721	WO 2005-FI4	20050104
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2004000005	A	20050706	FI 2004-5	20040105
FI 116141	B1	20050930		
CA 2551390	A1	20050721	CA 2005-2551390	20050104
EP 1704259	A1	20060927	EP 2005-701720	20050104
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS				
PRIORITY APPLN. INFO.:			FI 2004-5	A 20040105
			WO 2005-FI4	W 20050104

OTHER SOURCE(S): MARPAT 143:135160

AB Starch dissolved in an ionic liquid can be depolymd. without acid or base catalyst or enzyme. Starch is selectively depolymd. by mixing with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch by agitating at elevated temperature and for a period for time to effect depolymn. of the starch into desired depolymn. products. For example, all the starch was depolymd. into monomeric products by stirring a mixture of 150 mg of oven-dried native barley starch with 3 mL 1-butyl-3-methylimidazolium chloride solvent for 30 min at 85° and 2h at 150°. Stirring a similar mixture for 30 min at 85° and 2h at 100° gave a product mixture containing monomeric products of depolymd. amylose but amylopectin remained intact (GPC).

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Starch depolymerization in ionic liquid solvents

IN Myllymaeki, Vesa; Aksela, Reijo

AB Starch dissolved in an ionic liquid can be depolymd. without acid or base catalyst or enzyme. Starch is selectively depolymd. by mixing with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch by agitating at elevated temperature and for a.

ST starch depolymn ionic liq solvent; butylmethylimidazolium chloride solvent starch depolymn

IT Depolymerization

(selective; starch selective depolymn. in ionic liquid solvents)

IT Ionic liquids

(solvents; starch selective depolymn. in)

IT Polysaccharides, processes

RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PROC (Process)

(starch selective depolymn. in ionic liquid solvents)
 IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
 RL: TEM (Technical or engineered material use); USES (Uses)
 (solvent; starch selective depolymn. in ionic liquid solvents)
 IT 9005-25-8, Starch, processes
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PROC (Process)
 (starch selective depolymn. in ionic liquid solvents)

L6 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:523500 CAPLUS
 DOCUMENT NUMBER: 143:28326
 TITLE: Etherification of cellulose in ionic liquid solutions
 INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo
 PATENT ASSIGNEE(S): Kemira Oyj, Finland
 SOURCE: PCT Int. Appl., 23 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005054298	A1	20050616	WO 2004-FI730	20041202
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2003001763	A	20050604	FI 2003-1763	20031203
FI 116140	B1	20050930		
CA 2548007	A1	20050616	CA 2004-2548007	20041202
EP 1689788	A1	20060816	EP 2004-801227	20041202
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS				
US 2007112185	A1	20070517	US 2007-581491	20070116
PRIORITY APPLN. INFO.:			FI 2003-1763	A 20031203
			WO 2004-FI730	W 20041202

OTHER SOURCE(S): MARPAT 143:28326

AB Cellulose is mixed and dissolved in an ionic liquid solvent and the solution is treated with an etherifying agent in the presence of inorg. base to form a cellulose ether, which is subsequently separated from the solution

The dissoln. and the etherification are carried out in the absence of organic base and in the substantial absence of H₂O. Microwave irradiation and/or pressure can be applied to assist in dissoln. and etherification. Thus, 50 mg cellulose was dissolved in 5 g 1-butyl-3-methylimidazolium chloride (m. 60°) with the aid of microwaves to give 1% solution ClCH₂CO₂H (2.05 equiv) was added to the solution followed by 3.25 equiv of solid NaOH,

the reaction mixture was heated for 2 h at 100° under microwave radiation and the resulting CM-cellulose was precipitated with MeOH, washed with MeOH and 80% aqueous MeOH, and dried.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Etherification of cellulose in ionic liquid solutions
IN Myllymaeki, Vesa; Aksela, Reijo
AB Cellulose is mixed and dissolved in an ionic liquid solvent and the solution is treated with an etherifying agent in the presence of inorg. base to form a. . .
ST cellulose etherification ionic liq solvent microwave; butylmethylimidazolium chloride solvent CM cellulose manuf; chloroacetic acid etherification cellulose butylmethylimidazolium chloride solvent
IT Etherification
Ionic liquids
(etherification of cellulose in ionic liquid solution)
IT Microwave
(etherification of cellulose in ionic liquid solution in presence of)
IT 9004-32-4P, CM cellulose sodium salt
RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
(etherification of cellulose in ionic liquid solution)
IT 9004-34-6, Cellulose, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(etherification of cellulose in ionic liquid solution)
IT 79-11-8, Chloroacetic acid, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(etherification of cellulose; etherification of cellulose in ionic liquid solution)
IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
RL: TEM (Technical or engineered material use); USES (Uses)
(solvent; etherification of cellulose in ionic liquid solution)

L6 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:239036 CAPLUS

DOCUMENT NUMBER: 142:299721

TITLE: Esterification of starch under microwave irradiation and pressure

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2005023873	A1	20050317	WO 2004-FI523	20040910
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,			

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG

FI 2003001301	A	20050312	FI 2003-1301	20030911
FI 116142	B1	20050930		
CA 2533553	A1	20050317	CA 2004-2533553	20040910
EP 1664125	A1	20060607	EP 2004-767037	20040910
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004013432	A	20061010	BR 2004-13432	20040910
US 2007073051	A1	20070329	US 2006-566975	20061207
PRIORITY APPLN. INFO.:			FI 2003-1301	A 20030911
			WO 2004-FI523	W 20040910

OTHER SOURCE(S): MARPAT 142:299721

AB An organic starch ester is prepared by mixing a starch material, such as natural starch or hydrolyzed starch, with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch with an organic esterifying agent, such as C1-11 carboxylic acid, to form an organic starch ester, and subsequently separating the organic starch ester from the solution by adding a non-solvent, such as alcs., ketones, and acetonitrile, to the starch ester solution. Microwave irradiation and/or pressure can be applied to assist the dissoln. and esterification. Thus, native barely starch was dissolved in ionic 1-butyl-3-methylimidazolium chloride and reacted with acetic anhydride, followed by quenching with ethanol to receive starch acetate.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

IN Myllymaeki, Vesa; Aksela, Reijo

AB An organic starch ester is prepared by mixing a starch material, such as natural starch or hydrolyzed starch, with an ionic liquid solvent to dissolve the starch, and then treating the dissolved starch with an organic esterifying agent, such as C1-11. . . Microwave irradiation and/or pressure can be applied to assist the dissoln. and esterification. Thus, native barely starch was dissolved in ionic 1-butyl-3-methylimidazolium chloride and reacted with acetic anhydride, followed by quenching with ethanol to receive starch acetate.

L6 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:158715 CAPLUS

DOCUMENT NUMBER: 142:242565

TITLE: Dissolution and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

-----	-----	-----	-----	-----
WO 2005017001	A1	20050224	WO 2004-FI476	20040813
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2003001156	A	20050216	FI 2003-1156	20030815
FI 115835	B1	20050729		
CA 2532989	A1	20050224	CA 2004-2532989	20040813
EP 1654307	A1	20060510	EP 2004-742219	20040813
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004013435	A	20061010	BR 2004-13435	20040813
PRIORITY APPLN. INFO.:			FI 2003-1156	A 20030815
			WO 2004-FI476	W 20040813

OTHER SOURCE(S): MARPAT 142:242565

AB Wood, straw, and other natural lignocellulosic materials can be dissolved in an ionic liquid solvent under microwave irradiation and/or under pressure, and cellulose and other organic compds., such as lignin and extractives, can also be separated from the solution by precipitating with non-solvent ,
such as water, alcs., ketones, and ethers, of cellulose. Thus, plywood sawdust was dissolved in 1-butyl-3-methyl-imidazolium chloride under microwave irradiation

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Dissolution and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation

IN Myllymaeki, Vesa; Aksela, Reijo

AB Wood, straw, and other natural lignocellulosic materials can be dissolved in an ionic liquid solvent under microwave irradiation and/or under pressure, and cellulose and other organic compds., such as lignin and extractives, can. . . .

ST dissoln delignification lignocellulosic ionic liq solvent
microwave irradsn; plywood sawdust wood straw butylmethylimidazolium chloride dissoln microwave irradsn

IT Wood
(chips; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Dissolution
Straw
Wood
(dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Solvents
(ionic, liquid; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Microwave
(irradiation; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Wood boards
 (plywood, sawdust; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Sawdust
 (plywood; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Wood
 (soft; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT Alcohols, uses
 Ethers, uses
 Ketones, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (solvent; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT 9004-34-6P, Cellulose, preparation 9005-53-2P, Lignin, preparation
 RL: PUR (Purification or recovery); PREP (Preparation)
 (dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

IT 79917-90-1, 1-Butyl-3-methyl-imidazolium chloride
 RL: NUU (Other use, unclassified); USES (Uses)
 (solvent; dissoln. and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation)

=> file stng

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	28.13	28.34
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-4.68	-4.68

FILE 'STNGUIDE' ENTERED AT 16:00:24 ON 26 NOV 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
 LAST RELOADED: Nov 23, 2007 (20071123/UP).

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.06	28.40
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-4.68

FILE 'CAPLUS' ENTERED AT 16:00:34 ON 26 NOV 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is

held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 26 Nov 2007 VOL 147 ISS 23
FILE LAST UPDATED: 25 Nov 2007 (20071125/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> e swatloski/au

E1	1	SWATLING D K/AU
E2	5	SWATLING DONALD K/AU
E3	0 -->	SWATLOSKE/AU
E4	1	SWATLOSKE R A/AU
E5	3	SWATLOSKE R P/AU
E6	84	SWATLOSKE RICHARD P/AU
E7	3	SWATLOSKE RICHARD PATRICK/AU
E8	2	SWATLOSKE ROBERT/AU
E9	2	SWATLOSKE ROBERT A/AU
E10	1	SWATMAN C C/AU
E11	2	SWATMAN DAVID R/AU
E12	1	SWATMAN LESLIE/AU

=> s e4-e7

	1	"SWATLOSKE R A"/AU
	3	"SWATLOSKE R P"/AU
	84	"SWATLOSKE RICHARD P"/AU
	3	"SWATLOSKE RICHARD PATRICK"/AU
L7	91	("SWATLOSKE R A"/AU OR "SWATLOSKE R P"/AU OR "SWATLOSKE RICHARD P"/AU OR "SWATLOSKE RICHARD PATRICK"/AU)

=> s 17 and "ionic liquid"

	284921	"IONIC"
	511	"IONICS"
	285185	"IONIC"
		("IONIC" OR "IONICS")
	800777	"LIQUID"
	138530	"LIQUIDS"
	904338	"LIQUID"
		("LIQUID" OR "LIQUIDS")
	1105042	"LIQ"
	104871	"LIQS"
	1145172	"LIQ"
		("LIQ" OR "LIQS")
	1588658	"LIQUID"
		("LIQUID" OR "LIQ")
	11079	"IONIC LIQUID"
		("IONIC" (W) "LIQUID")
L8	83	L7 AND "IONIC LIQUID"

=> s 18 and cellulose
360710 CELLULOSE
4428 CELLULOSES
361213 CELLULOSE
(CELLULOSE OR CELLULOSES)
L9 26 L8 AND CELLULOSE

=> d 19 1-26 ibib kwic

L9 ANSWER 1 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:681891 CAPLUS

DOCUMENT NUMBER: 147:202400

TITLE: Sensor technologies based on a cellulose
supported platform

AUTHOR(S): Poplin, Jane Holly; Swatloski, Richard P.;
Holbrey, John D.; Spear, Scott K.; Metlen, Andreas;
Gratzel, Michael; Nazeeruddin, Mohammad K.; Rogers,
Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Chemical Communications (Cambridge, United Kingdom)
(2007), (20), 2025-2027
CODEN: CHCOFS; ISSN: 1359-7345

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Sensor technologies based on a cellulose supported platform

AU Poplin, Jane Holly; Swatloski, Richard P.; Holbrey, John D.;
Spear, Scott K.; Metlen, Andreas; Gratzel, Michael; Nazeeruddin, Mohammad
K.; Rogers, Robin D.

AB A simple approach to sensor development based on encapsulating a probe
mol. in a cellulose support followed by regeneration from an
ionic liquid solution is demonstrated here by the codissoln.
of cellulose and 1-(2-pyridylazo)-2-naphthol in
1-butyl-3-methylimidazolium chloride followed by regeneration with water
to form strips which exhibit a proportionate (1: 1) response. . .

ST sensor technol cellulose supported platform

IT Colorimetric indicators

Spectrophotometry

(mercury determination in water by spectrophotometry with indicator-
cellulose composite)

IT Encapsulation

Ionic liquids

Polymer-supported reagents

(sensor technologies based on cellulose supported platform
for encapsulation of indicators)

IT 7732-18-5, Water, analysis

RL: AMX (Analytical matrix); ANST (Analytical study)

(mercury determination in water by spectrophotometry with indicator-
cellulose composite)

IT 7439-97-6, Mercury, analysis

RL: ANT (Analyte); ANST (Analytical study)

(mercury determination in water by spectrophotometry with indicator-

- cellulose composite)
- IT 85-85-8, 1-(2-Pyridylazo)-2-naphthol 403790-50-1
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (mercury determination in water by spectrophotometry with indicator-
cellulose composite)
- IT 9004-34-6, Cellulose, uses
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (sensor technologies based on cellulose supported platform
 for encapsulation of indicators)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
 RL: ARU (Analytical role, unclassified); ANST (Analytical study)
 (sensor technologies based on cellulose supported platform
 for encapsulation of indicators)

L9 ANSWER 2 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:10320 CAPLUS

DOCUMENT NUMBER: 147:450565

TITLE: Can ionic liquids dissolve wood?
 Processing and analysis of lignocellulosic materials
 with 1-n-butyl-3-methylimidazolium chloride

AUTHOR(S): Fort, Diego A.; Remsing, Richard C.; Swatloski,
Richard P.; Moyna, Patrick; Moyna, Guillermo;
 Rogers, Robin D.

CORPORATE SOURCE: Facultad de Quimica, Universidad de la Republica,
 Montevideo, 11800, Urug.

SOURCE: Green Chemistry (2007), 9(1), 63-69
 CODEN: GRCHFJ; ISSN: 1463-9262

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Can ionic liquids dissolve wood? Processing and
 analysis of lignocellulosic materials with 1-n-butyl-3-methylimidazolium
 chloride

AU Fort, Diego A.; Remsing, Richard C.; Swatloski, Richard P.;
 Moyna, Patrick; Moyna, Guillermo; Rogers, Robin D.

AB The bulk of the cellulose currently employed by industry is
 isolated from wood through Kraft pulping, a process which traditionally
 involves a barrage of environmentally. . . novel alternative approach
 for the processing of lignocellulosic materials that relies on their solubility
 in solvent systems based on the ionic liquid (IL)
 1-n-butyl-3-methylimidazolium chloride ([C4mim]Cl). Dissoln. profiles for
 woods of different hardness are presented, making emphasis on the direct
 anal. of. . . material and lignin content in the resulting liquors by
 means of conventional ¹³C NMR techniques. The authors also show that
cellulose can be readily reconstituted from the IL-based wood
 liquors in fair yields by the addition of a variety of precipitating. . .

in this

manner is virtually free of lignin and hemicellulose and has
 characteristics that are comparable to those of pure cellulose
 samples subjected to similar processing conditions.

ST wood lignocellulose dissoln butylmethylimidazolium chloride ionic
lig

IT Wood

(eucalyptus; processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT Wood
 (oak; processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT Wood
 (pine; processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT Wood
 (poplar; processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT Ionic liquids
 Pulping
 (processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT 75-05-8, Acetonitrile, uses 75-09-2, Dichloromethane, uses 9004-34-6,
Cellulose, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT 11132-73-3, Lignocellulose
 RL: PRP (Properties)
 (processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

IT 9005-53-2, Lignin, processes 9034-32-6, Hemicellulose
 RL: REM (Removal or disposal); PROC (Process)
 (processing of lignocellulosic materials by dissoln. in
 (butyl)methylimidazolium chloride ionic liquid)

L9 ANSWER 3 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1215073 CAPLUS

DOCUMENT NUMBER: 147:450541

TITLE: Preparation of magnetic cellulose composites
 using ionic liquids

AUTHOR(S): Swatloski, Richard P.; Holbrey, John D.;
 Weston, James L.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry, Center for Green
 Manufacturing, The University of Alabama, Tuscaloosa,
 AL, 35487, USA

SOURCE: Chimica Oggi (2006), 24(2), 31-32, 34-35
 CODEN: CHOGDS; ISSN: 0392-839X

PUBLISHER: Tekno Scienze

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Preparation of magnetic cellulose composites using ionic
liquids

AU Swatloski, Richard P.; Holbrey, John D.; Weston, James L.;
 Rogers, Robin D.

AB Cellulose-magnetite composites have been prepared by suspension
 and dispersion of magnetite particles in a homogeneous ionic
liquid solution of cellulose, followed by regeneration into
 water, enabling the preparation of magnetically responsive films, floes,
 fibers, or beads. The materials prepared were. . .

ST magnetic cellulose composite ionic liq prepn
 property

IT Dispersion (of materials)
 Encapsulation

Ionic liquids

Magnetic materials

Particle size

Polymer morphology

Remanence

Suspensions

Thermal stability

(preparation of magnetic cellulose composites using ionic liqs. by suspension and dispersion of magnetite particles and characterization of composites)

IT Paramagnetism

(superparamagnetism; preparation of magnetic cellulose composites using ionic liqs. by suspension and dispersion of magnetite particles and characterization of composites)

IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride

RL: TEM (Technical or engineered material use); USES (Uses)

(ionic liquid; preparation of magnetic cellulose composites using ionic liqs. by suspension and dispersion of magnetite particles and characterization of composites)

IT 1309-38-2, Magnetite, uses

RL: MOA (Modifier or additive use); PEP (Physical, engineering or chemical process); PROC (Process); USES (Uses)

(preparation of magnetic cellulose composites using ionic liqs. by suspension and dispersion of magnetite particles and characterization of composites)

IT 9004-34-6, Cellulose, uses

RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PRP (Properties); PROC (Process); USES (Uses)

(preparation of magnetic cellulose composites using ionic liqs. by suspension and dispersion of magnetite particles and characterization of composites)

L9 ANSWER 4 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1187097 CAPLUS

DOCUMENT NUMBER: 147:236580

TITLE: Keynote address: new solvent for cellulose extrusion

AUTHOR(S): Broughton, Roy; Wang, Weijun; Shen, Guanglin; Farag, Ramsis; Swatloski, Richard P.; Rogers, Robin D.

CORPORATE SOURCE: Auburn University, Auburn, AL, USA

SOURCE: Proceedings - Beltwide Cotton Conferences (2005) 3291/1-3291/5

CODEN: PCOCEN; ISSN: 1059-2644

PUBLISHER: National Cotton Council

DOCUMENT TYPE: Journal; (computer optical disk)

LANGUAGE: English

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Keynote address: new solvent for cellulose extrusion

AU Broughton, Roy; Wang, Weijun; Shen, Guanglin; Farag, Ramsis; Swatloski, Richard P.; Rogers, Robin D.

AB A variety of celluloses have been dissolved in the ionic liquid 1-butyl-3-methylimidazolium chloride. The solns. were extruded in a dry-jet, wet-spinning process using water as a coagulation bath to produce a . . . having a tenacity of 2.0-4.4 g/denier (1.8-4.0 cN/dtex) and a breaking elongation of 4-20% depending on the extrusion

conditions. This ionic liquid appears to be versatile as a cellulose extrusion solvent with minimal polymer degradation. As none of the extrusion, coagulation, and drawing conditions have been optimized, the authors conclude that this new solvent has significant potential for the manufacture of regenerated cellulose fibers.

ST ionic liq solvent extrusion regenerated cellulose fiber; butylmethylimidazolium chloride solvent spinning regenerated cellulose fiber

IT Cellulose pulp
Cotton fibers
(ionic liquid solvent for extrusion of regenerated cellulose fibers from)

IT Elongation at break
Tenacity
Young's modulus
(of regenerated cellulose fibers spun from ionic liquid solvent)

IT Ionic liquids
(solvents for extrusion of regenerated cellulose fibers)

IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
RL: NUU (Other use, unclassified); USES (Uses)
(ionic liquid solvent for extrusion of regenerated cellulose fibers)

L9 : ANSWER 5 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1158863 CAPLUS

DOCUMENT NUMBER: 147:119949

TITLE: Effects of MAPP as coupling agent on the performance of regenerated cellulose film reinforced polypropylene composites

AUTHOR(S): Haque, A.; Mobley, C.; Daly, D. T.; Rogers, R. D.; Swatloski, R. P.; Ramasetty, A.

CORPORATE SOURCE: Department of Aerospace Engineering and Mechanics, The University of Alabama, Tuscaloosa, AL, 35487, USA

SOURCE: Proceedings of the American Society for Composites, Technical Conference (2006), 21st, 272/1-272/15
CODEN: PAMTEG; ISSN: 1084-7243

PUBLISHER: DEStech Publications, Inc.

DOCUMENT TYPE: Journal; (computer optical disk)

LANGUAGE: English

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Effects of MAPP as coupling agent on the performance of regenerated cellulose film reinforced polypropylene composites

AU Haque, A.; Mobley, C.; Daly, D. T.; Rogers, R. D.; Swatloski, R. P.; Ramasetty, A.

AB In this paper regenerated cellulose films were investigated for possible reinforcement in thermoplastic polypropylene resin. The pulp-based cellulose films were processed using ionic liqs. (ILs) with maleic anhydride polypropylene (MAPP) as a compatibilizer. These regenerated compatible cellulose films were incorporated in polypropylene resin to process bio-polymer matrix laminate composites. To improve compatibility of polypropylene resin and cellulose film, various weight percentages of MAPP binder were incorporated into cellulose film to achieve strong bonding and optimum stress transfer at the interface. Initially, the mech. properties of cellulose (CE) film and neat polypropylene (PP1, PP2) resins

were determined as baseline data. The effects of various MAPP concns. on tensile, flexural, and interlaminar shear properties, moisture absorption and m.p. of cellulose/polypropylene (CE/PP) composites were investigated. Microstructural examns. were conducted using optical and SEM to study the structure and fracture surface of. . . Reasonable agreements were observed between the exptl. and theor. predicted data for well bonded specimens. The fracture anal. showed that cellulose film with 22% MAPP concentration provides excellent adhesion between neat PP resin and CE film reinforcement. Improved adhesion between CE. . .

ST polypropylene cellulose laminated composite prepn property;
cellulose film polypropylene laminated composite maleated polypropylene coupler

IT Cellophane
Coupling agents
Elongation at break
Melting point
Stress-strain relationship
Tensile strength
Young's modulus
(effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT Laminated plastics, uses
RL: PRP (Properties); TEM (Technical or engineered material use); USES (Uses)
(effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT Polymer morphology
(fracture-surface; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT Absorption
(of water; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT Fracture surface morphology
(polymeric; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT Stress, mechanical
(yield; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT 7732-18-5, Water, properties
RL: PRP (Properties)
(absorption; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT 108-31-6D, Maleic anhydride, reaction products with polypropylene
9003-07-0D, Polypropylene, maleated
RL: MOA (Modifier or additive use); USES (Uses)
(coupler; effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

IT 9003-07-0, Polypropylene
RL: PRP (Properties); TEM (Technical or engineered material use); USES

(Uses)

(effects of maleated polypropylene as coupling agent on performance of regenerated cellulose film-reinforced laminated polypropylene composites)

L9 ANSWER 6 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:247215 CAPLUS

TITLE: How understanding the ionic liquid /cellulose dissolution mechanism can guide the generation of advanced cellulose-based materials

AUTHOR(S): Swatloski, Richard P.; Broughton, Roy M.; Moyna, Guillermo; Daly, Dan T.; Spear, Scott K.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green Manufacturing, The University of Alabama, Tuscaloosa, AL, 35487, USA

SOURCE: Abstracts of Papers, 231st ACS National Meeting, Atlanta, GA, United States, March 26-30, 2006 (2006), IEC-204. American Chemical Society: Washington, D. C. CODEN: 69HYEC

DOCUMENT TYPE: Conference; Meeting Abstract; (computer optical disk)

LANGUAGE: English

TI How understanding the ionic liquid/cellulose dissolution mechanism can guide the generation of advanced cellulose-based materials

AU Swatloski, Richard P.; Broughton, Roy M.; Moyna, Guillermo; Daly, Dan T.; Spear, Scott K.; Rogers, Robin D.

L9 ANSWER 7 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:247162 CAPLUS

TITLE: Use of ionic liquids for the processing and analysis of lignocellulosic materials

AUTHOR(S): Remsing, Richard C.; Fort, Diego A.; Swatloski, Richard P.; Moyna, Patrick; Rogers, Robin D.; Moyna, Guillermo

CORPORATE SOURCE: Department of Chemistry & Biochemistry, University of the Sciences in Philadelphia, Philadelphia, PA, 19104, USA

SOURCE: Abstracts of Papers, 231st ACS National Meeting, Atlanta, GA, United States, March 26-30, 2006 (2006), IEC-151. American Chemical Society: Washington, D. C. CODEN: 69HYEC

DOCUMENT TYPE: Conference; Meeting Abstract; (computer optical disk)

LANGUAGE: English

TI Use of ionic liquids for the processing and analysis of lignocellulosic materials

AU Remsing, Richard C.; Fort, Diego A.; Swatloski, Richard P.; Moyna, Patrick; Rogers, Robin D.; Moyna, Guillermo

AB Cellulose is the most abundant renewable biopolymer on Earth. While its most notable uses are related to the paper and textile. . . the Kraft pulping process using a barrage of environmentally detrimental chems. We describe a simple and novel method to extract cellulose from wood using "green" solvent systems based on the ionic liquid (IL) 1-n-butyl-3-methylimidazolium chloride ([C4mim]Cl). Extraction profiles for different woods are presented, making particular emphasis on the anal. of cellulose content in the IL-based wood

liquors by means of ^{13}C NMR techniques. In addition, we show that cellulose virtually free of lignin can be easily reconstituted from the IL liquors. Modifications to the methodol., including the use of. . .

L9 ANSWER 8 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:233428 CAPLUS

DOCUMENT NUMBER: 144:452142

TITLE: Mechanism of cellulose dissolution in the ionic liquid 1-n-butyl-3-methylimidazolium chloride: a ^{13}C and $^{35/37}\text{Cl}$ NMR relaxation study on model systems

AUTHOR(S): Remsing, Richard C.; Swatloski, Richard P.; Rogers, Robin D.; Moyna, Guillermo

CORPORATE SOURCE: Department of Chemistry & Biochemistry, University of the Sciences in Philadelphia, Philadelphia, PA, 19104, USA

SOURCE: Chemical Communications (Cambridge, United Kingdom) (2006), (12), 1271-1273

CODEN: CHCOFS; ISSN: 1359-7345

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- TI Mechanism of cellulose dissolution in the ionic liquid 1-n-butyl-3-methylimidazolium chloride: a ^{13}C and $^{35/37}\text{Cl}$ NMR relaxation study on model systems
- AU Remsing, Richard C.; Swatloski, Richard P.; Rogers, Robin D.; Moyna, Guillermo
- AB ^{13}C and $^{35/37}\text{Cl}$ NMR relaxation measurements on several model systems demonstrate that the solvation of cellulose by the ionic liquid (IL) 1-n-butyl-3-methylimidazolium chloride ($[\text{C4mim}]\text{Cl}$) involves hydrogen bonding between the carbohydrate hydroxyl protons and the IL chloride ions in a 1:1. . .
- ST mechanism cellulose dissoln ionic liq butyl methylimidazolium chloride
- IT Dissolution
Hydrogen bond
(^{13}C and $^{35/37}\text{Cl}$ NMR relaxation study of mechanism of cellulose dissoln. in ionic liquid 1-n-butyl-3-methylimidazolium chloride)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
RL: NUU (Other use, unclassified); USES (Uses)
(ionic liquid; ^{13}C and $^{35/37}\text{Cl}$ NMR relaxation study of mechanism of cellulose dissoln. in ionic liquid 1-n-butyl-3-methylimidazolium chloride)
- IT 50-99-7, Glucose, processes 528-50-7, D-Cellobiose 604-69-3, β -D-Glucose pentaacetate 9004-34-6, Cellulose, processes
RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)
(^{13}C and $^{35/37}\text{Cl}$ NMR relaxation study of mechanism of cellulose dissoln. in ionic liquid 1-n-butyl-3-methylimidazolium chloride)

L9 ANSWER 9 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:1130915 CAPLUS

DOCUMENT NUMBER: 143:387946
 TITLE: Polymer dissolution and blend formation in
ionic liquids
 INVENTOR(S): Holbrey, John D.; Swatloski, Richard P.;
 Chen, Ji; Daly, Dan; Rogers, Robin D.
 PATENT ASSIGNEE(S): The University of Alabama, USA; Holbrey, John, D.;
 Swatloski, Richard, P.; Rogers, Robin, D.
 SOURCE: PCT Int. Appl., 32 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005098546	A2	20051020	WO 2005-US10235	20050325
WO 2005098546	A3	20070222		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2005288484	A1	20051229	US 2005-87496	20050324
AU 2005231083	A1	20051020	AU 2005-231083	20050325
CA 2560680	A1	20051020	CA 2005-2560680	20050325
EP 1733282	A2	20061220	EP 2005-729932	20050325
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, LV, MK, YU				
BR 2005009250	A	20070911	BR 2005-9250	20050325
JP 2007530743	T	20071101	JP 2007-505253	20050325
IN 2006DN05591	A	20070824	IN 2006-DN5591	20060925
NO 2006004827	A	20061025	NO 2006-4827	20061025
KR 2007042118	A	20070420	KR 2006-722385	20061026
PRIORITY APPLN. INFO.:			US 2004-556484P	P 20040326
			WO 2005-US10235	W 20050325

TI Polymer dissolution and blend formation in ionic liquids
 IN Holbrey, John D.; Swatloski, Richard P.; Chen, Ji; Daly, Dan;
 Rogers, Robin D.

AB The ionic liqs. are for the dissoln. of various
 polymers and/or copolymers, the formation of resins and blends, and the
 reconstitution of polymer and/or copolymer solns., and the dissoln. and
 blending of functional additives and/or various polymers and/or
 copolymers. Thus, ≥ 1 ionic liquid, which is a
 liquid salt complex that exists in the liquid phase between about -70 to
 300°, is mixed with ≥ 2 differing polymeric materials to form
 a mixture, and adding a nonsolvent to the mixture to remove the ionic
liquid from the resin or blend.

ST ionic liq polymer blend cellulose
 IT Polyimides, uses

- RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PYP (Physical process); PROC (Process); USES (Uses)
(polyamide-; polymer blend formation in ionic liqs
.)
- IT Polyamides, uses
RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PYP (Physical process); PROC (Process); USES (Uses)
(polyimide-; polymer blend formation in ionic liqs
.)
- IT Polyamides, uses
Polyesters, uses
Polyimides, uses
Polyoxyalkylenes, uses
RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PYP (Physical process); PROC (Process); USES (Uses)
(polymer blend formation in ionic liqs.)
- IT Polymer blends
RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)
(polymer blend formation in ionic liqs.)
- IT Ionic liquids
(solvents; polymer blend formation in ionic liqs.)
- IT 25233-30-1, Polyaniline
RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PYP (Physical process); PROC (Process); USES (Uses)
(emeraldine base; polymer blend formation in ionic liqs.)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
RL: NUU (Other use, unclassified); USES (Uses)
(polymer blend formation in ionic liqs.)
- IT 9002-89-5, Polyvinyl alcohol 9002-98-6 9004-34-6, Cellulose,
uses 9005-25-8, Starch, uses 9034-32-6, Hemicellulose 25014-41-9,
Polyacrylonitrile 25249-16-5, Poly(2-hydroxyethyl methacrylate)
25322-68-3, Polyethylene glycol 26913-06-4, Poly[imino(1,2-ethanediy)]
RL: PEP (Physical, engineering or chemical process); POF (Polymer in formulation); PYP (Physical process); PROC (Process); USES (Uses)
(polymer blend formation in ionic liqs.)

L9 ANSWER 10 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:734798 CAPLUS

DOCUMENT NUMBER: 144:332128

TITLE: Applying ionic liquids for
controlled processing of polymer materials

AUTHOR(S): Holbrey, John D.; Chen, Ji; Turner, Megan B.;
Swatloski, Richard P.; Spear, Scott K.;
Rogers, Robin D.

CORPORATE SOURCE: Center for Green Manufacturing and Department of
Chemistry, The University of Alabama, Tuscaloosa, AL,
35487, USA

SOURCE: ACS Symposium Series (2005), 913(Ionic Liquids in
Polymer Systems), 71-87
CODEN: ACSMC8; ISSN: 0097-6156

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

REFERENCE COUNT: 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Applying ionic liquids for controlled processing of
polymer materials
AU Holbrey, John D.; Chen, Ji; Turner, Megan B.; Swatloski, Richard
P.; Spear, Scott K.; Rogers, Robin D.
AB A review. This perspective examines the potential, highlighting some
examples from the on-going research program, to evaluate and apply
ionic liqs. as advanced functional solvents for
dissolving and processing polymers to prepare active materials and
composites for sensor and smart materials. . .
ST review ionic liq solvent polymer processing composite
IT Green chemistry

Ionic liquids

Solvents

(applying ionic liqs. for controlled processing of
polymer materials in manufacture of composites)

IT Polymers, processes

RL: PEP (Physical, engineering or chemical process); PYP (Physical
process); PROC (Process)

(applying ionic liqs. for controlled processing of
polymer materials in manufacture of composites)

IT 9004-34-6, Cellulose, processes

RL: PEP (Physical, engineering or chemical process); PYP (Physical
process); PROC (Process)

(applying ionic liqs. for controlled processing of
polymer materials in manufacture of composites)

L9 ANSWER 11 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:342899 CAPLUS

DOCUMENT NUMBER: 143:28264

TITLE: High-resolution ¹³C NMR studies of cellulose
and cellulose oligomers in ionic
liquid solutions

AUTHOR(S): Moulthrop, Jason S.; Swatloski, Richard P.;
Moyna, Guillermo; Rogers, Robin D.

CORPORATE SOURCE: Dep. of Chem. and Biochem., Univ. of the Sci. in
Philadelphia, Philadelphia, PA, 19104, USA

SOURCE: Chemical Communications (Cambridge, United Kingdom)
(2005), (12), 1557-1559

CODEN: CHCOFS; ISSN: 1359-7345

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI High-resolution ¹³C NMR studies of cellulose and
cellulose oligomers in ionic liquid solutions

AU Moulthrop, Jason S.; Swatloski, Richard P.; Moyna, Guillermo;
Rogers, Robin D.

AB High-resolution ¹³C NMR studies of cellulose and cellulose
oligomers dissolved in the ionic liquid (IL)
1-butyl-3-methylimidazolium chloride ([C4mim]Cl) show that the
β-(1→4)-linked glucose oligomers are disordered in this medium
and have a conformational behavior. . .

ST cellulose oligomer ionic liq NMR
conformation behavior

IT NMR spectroscopy

(carbon-13; conformational behavior of cellulose and

cellulose oligomers in ionic liquid solns.
studied by ¹³C NMR)

- IT Conformation
(β -; of cellulose and cellulose oligomers in
ionic liquid solns. studied by ¹³C NMR)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
RL: NUU (Other use, unclassified); USES (Uses)
(conformational behavior of cellulose and cellulose
oligomers in ionic liquid solns. studied by ¹³C NMR)
- IT 528-50-7, Cellobiose 2478-35-5, Cellohexaose 9004-34-6,
Cellulose, properties 38819-01-1, Cellotetraose
RL: PRP (Properties)
(conformational behavior of cellulose and cellulose
oligomers in ionic liquid solns. studied by ¹³C NMR)

L9 ANSWER 12 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:655919 CAPLUS

TITLE: High-resolution ¹³C NMR studies of amylose and
cellulose oligomers in 1-butyl-3-
methylimidazolium chloride solutions

AUTHOR(S): Moulthrop, Jason S.; Swatloski, Richard P.;
Rogers, Robin D.; Moyna, Guillermo

CORPORATE SOURCE: Department of Chemistry & Biochemistry, University of
the Sciences in Philadelphia, Philadelphia, PA, 19104,
USA

SOURCE: Abstracts of Papers, 228th ACS National Meeting,
Philadelphia, PA, United States, August 22-26, 2004
(2004), CARB-063. American Chemical Society:
Washington, D. C.
CODEN: 69FTZ8

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI High-resolution ¹³C NMR studies of amylose and cellulose
oligomers in 1-butyl-3-methylimidazolium chloride solutions

AU Moulthrop, Jason S.; Swatloski, Richard P.; Rogers, Robin D.;
Moyna, Guillermo

AB A high-resolution ¹³C NMR (NMR) spectroscopy study of amylose and
cellulose oligomers dissolved in the ionic liq
. (IL) 1-butyl-3-methylimidazolium chloride ([C4mim]Cl) is presented.
Results for all the oligosaccharides studied, which included linear
(1 \rightarrow 4)-linked glucose dimers, tetramers, and. . . angles, indicates
that the conformational preferences of these oligosaccharides in [C4mim]Cl
and aqueous solns. are similar. Preliminary results obtained for
cellulose show that its conformational behavior in [C4mim]Cl solution
parallels the one observed for the smaller β -(1 \rightarrow 4) glucose
oligomers, and that the polysaccharide is disordered in the IL solution The
impact that these findings may have on green cellulose
processing methods with potential industrial application is discussed.

L9 ANSWER 13 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:234276 CAPLUS

DOCUMENT NUMBER: 141:39087

TITLE: Applying ionic liquid solvent
characteristics for controlled processing of polymer
materials

AUTHOR(S): Holbrey, John D.; Chen, Ji; Turner, Megan B.;
Swatloski, Richard P.; Spear, Scott K.;

CORPORATE SOURCE: Rogers, Robin D.
Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA
SOURCE: Polymer Preprints (American Chemical Society, Division
of Polymer Chemistry) (2004), 45(1), 297-298
CODEN: ACPPAY; ISSN: 0032-3934
PUBLISHER: American Chemical Society, Division of Polymer
Chemistry
DOCUMENT TYPE: Journal; (computer optical disk)
LANGUAGE: English
REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- TI Applying ionic liquid solvent characteristics for
controlled processing of polymer materials
- AU Holbrey, John D.; Chen, Ji; Turner, Megan B.; Swatloski, Richard P.; Spear, Scott K.; Rogers, Robin D.
- AB The unique, and controllable solubility parameters exhibited by Ionic Liqs. (ILs) as a general class of fluids, and by individual IL examples, can be applied to polymer dissoln. and processing, . . .
- ST ionic liq polymer soly cellulose
- IT Ionic liquids
Polymer morphology
Solubility
(applying ionic liquid solvent characteristics for
controlled processing of polymer materials)
- IT DNA
Polyanilines
Polyoxyalkylenes, properties
RL: PRP (Properties)
(applying ionic liquid solvent characteristics for
controlled processing of polymer materials)
- IT Albumins, properties
RL: PRP (Properties)
(serum, bovine; applying ionic liquid solvent
characteristics for controlled processing of polymer materials)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride 174501-64-5,
1-Butyl-3-methylimidazolium hexafluorophosphate 174501-65-6,
1-Butyl-3-methylimidazolium tetrafluoroborate
RL: NUU (Other use, unclassified); USES (Uses)
(applying ionic liquid solvent characteristics for
controlled processing of polymer materials)
- IT 9002-89-5, Poly (vinyl alcohol) 9004-34-6, Cellulose,
properties 9005-25-8, Starch, properties 25014-41-9, Polyacrylonitrile
25233-30-1, Polyaniline 25249-16-5, Poly(2-hydroxyethylmethacrylate)
25322-68-3, Poly (ethylene glycol) 25322-69-4, Polypropylene glycol
RL: PRP (Properties)
(applying ionic liquid solvent characteristics for
controlled processing of polymer materials)

L9 ANSWER 14 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:222070 CAPLUS

TITLE: Cellulose-supported colorimetric sensors for
mercury ion detection

AUTHOR(S): Poplin, Jane Holly; Swatloski, Richard P.;
Holbrey, John D.; Spear, Scott K.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green

SOURCE: Manufacturing, The University of Alabama, Tuscaloosa, AL, 35486, USA
 Abstracts of Papers, 227th ACS National Meeting, Anaheim, CA, United States, March 28-April 1, 2004 (2004), CELL-024. American Chemical Society: Washington, D. C.
 CODEN: 69FGKM
 DOCUMENT TYPE: Conference; Meeting Abstract
 LANGUAGE: English
 TI Cellulose-supported colorimetric sensors for mercury ion detection
 AU Poplin, Jane Holly; Swatloski, Richard P.; Holbrey, John D.; Spear, Scott K.; Rogers, Robin D.
 AB Cellulose membranes are extremely porous and highly wettable which has advantages over hydrophobic, poorly wetting supports such as PVC or polyethylene for sensing in aqueous systems by providing fast transport of water-soluble ions to the active sensing sites. Responsive, colorimetric cellulose materials can be prepared by introducing a sensing moiety into cellulose-in-ionic liquid solns., providing a flexible route for forming indicating films by casting from water. PAN,1-(2-pyridylazo)-2-naphthol, a colorimetric complexant for transition metals can be readily incorporated into cellulose films, and is extremely responsive to metal-ions in solution, tuning from orange to a deep-red in the presence of mercury. . .
 L9 ANSWER 15 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:162249 CAPLUS
 DOCUMENT NUMBER: 140:201295
 TITLE: Regenerated cellulose matrix-encapsulated active substances and method therefor
 INVENTOR(S): Holbrey, John David; Spear, Scott K.; Turner, Megan B.; Swatloski, Richard Patrick; Rogers, Robin Don
 PATENT ASSIGNEE(S): The University of Alabama, USA; PG Research Foundation
 SOURCE: U.S. Pat. Appl. Publ., 21 pp., Cont.-in-part of U.S. Ser. No. 256,521.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004038031	A1	20040226	US 2003-394989	20030321
US 6808557	B2	20041026		
US 2003157351	A1	20030821	US 2002-256521	20020927
US 6824599	B2	20041130		
CN 101007853	A	20070801	CN 2007-10085298	20021003
AU 2004224375	A1	20041007	AU 2004-224375	20040319
CA 2519652	A1	20041007	CA 2004-2519652	20040319
WO 2004084627	A2	20041007	WO 2004-US8411	20040319
WO 2004084627	A3	20060105		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,

NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,
 TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
 BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,
 ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
 SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
 TD, TG

BR 2004008606	A	20060307	BR 2004-8606	20040319
EP 1648692	A2	20060426	EP 2004-757863	20040319
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
CN 1867448	A	20061122	CN 2004-80013560	20040319
JP 2006526673	T	20061124	JP 2006-507356	20040319
IN 2005DN04541	A	20070817	IN 2005-DN4541	20051006
PRIORITY APPLN. INFO.:				
			US 2001-326704P	P 20011003
			US 2002-256521	A2 20020927
			CN 2002-823875	A3 20021003
			US 2003-394989	A 20030321
			WO 2004-US8411	W 20040319

OTHER SOURCE(S): MARPAT 140:201295

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Regenerated cellulose matrix-encapsulated active substances and
 method therefor

IN Holbrey, John David; Spear, Scott K.; Turner, Megan B.; Swatloski,
Richard Patrick; Rogers, Robin Don

AB The process involves encapsulation or immobilization of the active solid
 substance in a cellulose framework by regenerating
cellulose dissolved in an ionic liquid solvent
 in a regenerating solution The active substance can be initially present in
 the ionic liquid or in the regenerating solvent either
 as a solution or dispersion. The invention is applicable to mol.
 encapsulation and to. . . to the formation of bulk materials with a
 wide range of morphol. forms. Thus, carbamoylmethylphosphine oxide (I)
 encapsulated in a cellulose matrix was realized by adding I to a
 10% solution of cellulose in 1-butyl-3-methylimidazolium chloride (
ionic liquid) under vigorous stirring and then removing
 the ionic liquid with water.

ST ionic liq regenerated cellulose
 encapsulation active substance; carbamoylmethylphosphine oxide
 encapsulation regenerated cellulose

IT Quaternary ammonium compounds, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (ionic liquid; regenerated cellulose
 matrix-encapsulated active substances and method therefor)

IT Encapsulation
 (regenerated cellulose matrix-encapsulated active substances
 and method therefor)

IT Ubiquinones
 RL: MSC (Miscellaneous)
 (regenerated cellulose matrix-encapsulated active substances
 and method therefor)

IT Albumins, miscellaneous
 RL: MSC (Miscellaneous)
 (serum, bovine; regenerated cellulose matrix-encapsulated
 active substances and method therefor)

IT 2580-56-5, Victoria blue B

RL: MSC (Miscellaneous)
 (dye; regenerated cellulose matrix-encapsulated active substances and method therefor)

IT 9004-34-6, Cellulose, uses 76296-24-7, Cellulose azure
 RL: TEM (Technical or engineered material use); USES (Uses)
 (encapsulation agent; regenerated cellulose matrix-encapsulated active substances and method therefor)

IT 83242-95-9, CMPO
 RL: MSC (Miscellaneous)
 (extractant; regenerated cellulose matrix-encapsulated active substances and method therefor)

IT 79917-90-1, 1-Butyl-3-methylimidazolium Chloride
 RL: NUU (Other use, unclassified); USES (Uses)
 (ionic liquid; regenerated cellulose matrix-encapsulated active substances and method therefor)

IT 553-12-8, Protoporphyrin IX 80498-15-3, Laccase
 RL: MSC (Miscellaneous)
 (regenerated cellulose matrix-encapsulated active substances and method therefor)

L9 ANSWER 16 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2003:684181 CAPLUS
 DOCUMENT NUMBER: 140:147858
 TITLE: Ionic liquids for the dissolution and regeneration of cellulose
 AUTHOR(S): Swatloski, Richard P.; Holbrey, John D.; Spear, Scott K.; Rogers, Robin D.
 CORPORATE SOURCE: Department of Chemistry and Center for Green Manufacturing, The University of Alabama, Tuscaloosa, AL, 35487-0336, USA
 SOURCE: Proceedings - Electrochemical Society (2002), 2002-19(Molten Salts XIII), 155-164
 CODEN: PESODO; ISSN: 0161-6374
 PUBLISHER: Electrochemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Ionic liquids for the dissolution and regeneration of cellulose
 AU Swatloski, Richard P.; Holbrey, John D.; Spear, Scott K.; Rogers, Robin D.
 AB There is an increasing willingness to develop new cellulose-based materials, particularly from homogeneous solution, due to the fact that cellulose is the earth's most abundant biorenewable resource. The opportunity to use ionic liqs. as solvents for clean extraction and processing of cellulose was investigated. Cellulose can be dissolved in a number of ionic liqs. and easily regenerated by contacting with water. This allows a simple, benign system for the processing of cellulose into fibers, monoliths, and films by forming into an aqueous phase. This has potential environmental and cost advantages over current.

ST alkylmethylimidazolium ionic liq dissoln cellulose rayon regeneration
 IT Ionic liquids

Thermal decomposition

(alkylmethylimidazolium-based ionic liqs. for
dissoln. and regeneration of cellulose)

IT Rayon, properties

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
(alkylmethylimidazolium-based ionic liqs. for
dissoln. and regeneration of cellulose)

IT 64697-40-1, 1-Methyl-3-octylimidazolium chloride 79917-90-1,
1-Butyl-3-methylimidazolium chloride 85100-77-2, 1-Butyl-3-
methylimidazolium bromide 171058-17-6, 1-Hexyl-3-methylimidazolium
chloride 174501-64-5, 1-Butyl-3-methylimidazolium hexafluorophosp[hate
174501-65-6, 1-Butyl-3-methylimidazolium tetrafluoroborate 344790-87-0,
1-Butyl-3-methylimidazolium thiocyanate

RL: NUU (Other use, unclassified); USES (Uses)
(alkylmethylimidazolium-based ionic liqs. for
dissoln. and regeneration of cellulose)

IT 9004-34-6, Cellulose, processes

RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP
(Physical process); PROC (Process)
(alkylmethylimidazolium-based ionic liqs. for
dissoln. and regeneration of cellulose)

L9 ANSWER 17 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:635072 CAPLUS

TITLE: Application of ionic liquid
technologies to nuclear separations

AUTHOR(S): Rogers, Robin D.; Holbrey, John D.; Spear, Scott K.;
Gutowski, Keith E.; Bridges, Nicholas J.; Cocalia,
Violina A.; Swatloski, Richard P.

CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Abstracts of Papers, 226th ACS National Meeting, New
York, NY, United States, September 7-11, 2003 (2003),
NUCL-092. American Chemical Society: Washington, D.
C.

CODEN: 69EKY9

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Application of ionic liquid technologies to nuclear
separations

AU Rogers, Robin D.; Holbrey, John D.; Spear, Scott K.; Gutowski, Keith E.;
Bridges, Nicholas J.; Cocalia, Violina A.; Swatloski, Richard P.

AB Room temperature Ionic Liqs. (ILs), organic salts that are liquid
at, or close to room temperature have great potential application for uses in
liquid-liquid. . . into an IL; immobilizing IL extractant phases on solid
supports; and utilization of the solubilizing power of ILs to prepare
cellulose-based materials for f-element sepns. ILs can thus be
considered as a new class of materials for nuclear sepns., distinct from.

L9 ANSWER 18 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:632691 CAPLUS

TITLE: Cellulose films regenerated from ILs and
their role as scaffolding for enzyme attachment via
glutaraldehyde

AUTHOR(S): Turner, Megan B.; Spear, Scott K.; Swatloski,

CORPORATE SOURCE: Richard P.; Holbrey, John D.; Rogers, Robin D.
Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Abstracts of Papers, 226th ACS National Meeting, New
York, NY, United States, September 7-11, 2003 (2003),
IEC-190. American Chemical Society: Washington, D. C.
CODEN: 69EKY9

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Cellulose films regenerated from ILs and their role as
scaffolding for enzyme attachment via glutaraldehyde

AU Turner, Megan B.; Spear, Scott K.; Swatloski, Richard P.;
Holbrey, John D.; Rogers, Robin D.

AB . . . enzyme stability and thus activity, and also enable simple
recovery and recycling of catalysts. Here we describe the investigation
of cellulose films containing encapsulated laccase, from *Rhus*
vernificera, prepared by dissoln. and casting from 1-butyl-3-methylimidazole
chloride ionic liquid solution. Laccase can be successfully
immobilized in cellulose films prepared in this way, while
maintaining enzymic activity, and the activity can be controlled by
pretreatment and processing methodologies.. . .

L9 ANSWER 19 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:632574 CAPLUS

TITLE: Ionic liquids as green solvents:

Engineering new bio-based materials

AUTHOR(S): Swatloski, Richard P.; Holbrey, John D.;
Spear, Scott K.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Abstracts of Papers, 226th ACS National Meeting, New
York, NY, United States, September 7-11, 2003 (2003),
IEC-090. American Chemical Society: Washington, D. C.
CODEN: 69EKY9

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Ionic liquids as green solvents: Engineering new
bio-based materials

AU Swatloski, Richard P.; Holbrey, John D.; Spear, Scott K.;
Rogers, Robin D.

AB . . . complement, or replace diminishing petroleum-based feed-stocks.
We have recently utilized 1-butyl-3-methylimidazolium chloride for the
dissoln. of nature's most abundant renewable resources--cellulose
. Because ionic liqs. can dissolve a wide range of
materials, it can be anticipated that they will offer a route for
incorporation of many functional mols. for sensing, recognition, or mol.
binding into modified cellulose materials that have not been
accessible in other traditional cellulose solvent systems. In
this presentation we will examine the phys. properties of these new
materials, as well as their possible. . .

L9 ANSWER 20 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:632439 CAPLUS

TITLE: CMPO-impregnated cellulosic materials from
ionic liquids for f-element

separations
AUTHOR(S): Rogers, Robin D.; Holbrey, John D.; Spear, Scott K.;
Gutowski, Keith E.; Swatloski, Richard P.
CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA
SOURCE: Abstracts of Papers, 226th ACS National Meeting, New
York, NY, United States, September 7-11, 2003 (2003),
IEC-045. American Chemical Society: Washington, D. C.
CODEN: 69EKY9
DOCUMENT TYPE: Conference; Meeting Abstract
LANGUAGE: English
TI CMPO-impregnated cellulosic materials from ionic liquids
for f-element separations
AU Rogers, Robin D.; Holbrey, John D.; Spear, Scott K.; Gutowski, Keith E.;
Swatloski, Richard P.
AB By taking advantage of the solubility of both cellulose and CMPO
(octyl(phenyl)-N,N-diisobutylcarbamoylmethyl phosphine oxide) in the
ionic liquid 1-butyl-3-methylimidazolium chloride, we have
prepared CMPO-impregnated cellulosic materials as flocs, beads, rods, and
membranes. Americium-241, plutonium-239, and uranium-233 all exhibit
significant partitioning from aqueous solns. to the cellulose
impregnated materials with increasing concns. of nitric acid. In this
presentation we will examine the phys. properties of these new. . .

L9 ANSWER 21 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:282640 CAPLUS

DOCUMENT NUMBER: 138:289216

TITLE: Dissolution and processing of cellulose
using ionic liquids,
cellulose solution, and regenerating
cellulose

INVENTOR(S): Swatloski, Richard Patrick; Rogers, Robin
Don; Holbrey, John David

PATENT ASSIGNEE(S): The University of Alabama, USA; Pg Research
Foundation, Inc.

SOURCE: PCT Int. Appl., 59 pp.
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003029329	A2	20030410	WO 2002-US31404	20021003
WO 2003029329	A3	20030731		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

CA 2462460	A1	20030410	CA 2002-2462460	20021003
AU 2002347788	A1	20030414	AU 2002-347788	20021003
EP 1458805	A2	20040922	EP 2002-784000	20021003
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2005506401	T	20050303	JP 2003-532567	20021003
CN 1596282	A	20050316	CN 2002-823875	20021003
NZ 532076	A	20050930	NZ 2002-532076	20021003
BR 2002013106	A	20060523	BR 2002-13106	20021003
CN 101007853	A	20070801	CN 2007-10085298	20021003
MX 2004PA03029	A	20050620	MX 2004-PA3029	20040331
ZA 2004002610	A	20041223	ZA 2004-2610	20040401
NO 2004001774	A	20040430	NO 2004-1774	20040430
IN 2004DN01175	A	20060728	IN 2004-DN1175	20040430
PRIORITY APPLN. INFO.:			US 2001-326704P	P 20011003
			CN 2002-823875	A3 20021003
			WO 2002-US31404	W 20021003

OTHER SOURCE(S): MARPAT 138:289216

TI Dissolution and processing of cellulose using ionic liquids, cellulose solution, and regenerating cellulose

IN Swatloski, Richard Patrick; Rogers, Robin Don; Holbrey, John David

AB Cellulose is dissolved in an ionic liquid without derivatization, and is regenerated in a range of structural forms without requiring the use of harmful or volatile organic solvents. Cellulose solubility and the solution properties can be controlled by the selection of the ionic liquid constituents, with small cations and halide or pseudohalide anions favoring solution; dissoln. can be aided by irradiation. An ionic liquid, [C4mim]Cl, proved to be the best for dissolving cellulose.

ST cellulose dissolving ionic liq

IT Fibers

RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); PROC (Process)

(cellulosic; dissoln. and processing of cellulose using hydrophilic ionic liqs.)

IT Cellulose pulp

Dissolution

Gossypium hirsutum

Ionic liquids

Paper

Solvent effect

(dissoln. and processing of cellulose using hydrophilic ionic liqs.)

IT Crystallinity

(liquid; dissoln. and processing of cellulose using hydrophilic ionic liqs.)

IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride 85100-77-2, 1-Butyl-3-methylimidazolium bromide 344790-87-0, 1-Butyl-3-methylimidazolium thiocyanate

RL: NUU (Other use, unclassified); USES (Uses)

(dissoln. and processing of cellulose using hydrophilic ionic liqs.)

IT 9004-34-6, Cellulose, processes

RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); PROC (Process)

(dissoln. and processing of cellulose using hydrophilic
ionic liqs.)

L9 ANSWER 22 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:182947 CAPLUS

TITLE: Properties of regenerated cellulose from
ionic liquids

AUTHOR(S): Swatloski, Richard P.; Holbrey, John D.;
Spear, Scott K.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Abstracts of Papers, 225th ACS National Meeting, New
Orleans, LA, United States, March 23-27, 2003 (2003),
IEC-167. American Chemical Society: Washington, D. C.
CODEN: 69DSA4

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Properties of regenerated cellulose from ionic
liquids

AU Swatloski, Richard P.; Holbrey, John D.; Spear, Scott K.;
Rogers, Robin D.

AB Cellulose is the earth's most abundant biorenewable material.
It has important com. applications across a wide variety of technologies.
A hindrance in this field is due to the limited number of solvents capable of
completely dissolving cellulose. In our labs., we have utilized
ionic liqs. (ILs) to successfully dissolve up to 30%
wt/wt of cellulose without pretreatment or derivitization.
Cellulose can easily be regenerated from the IL solution simply by
contacting the cellulosic solution with water. We will describe methods for
regeneration-capable of forming thin films, filaments, and membranes.
Results from regenerated cellulose, as well as important phys.
properties of the regenerated cellulose will be discussed. This
research is sponsored by The PG Research Foundation, Inc.

L9 ANSWER 23 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:179636 CAPLUS

TITLE: Ionic liquids as green solvents
for the dissolution and regeneration of
cellulose

AUTHOR(S): Swatloski, Richard P.; Spear, Scott K.;
Holbrey, John D.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA

SOURCE: Abstracts of Papers, 225th ACS National Meeting, New
Orleans, LA, United States, March 23-27, 2003 (2003),
CELL-131. American Chemical Society: Washington, D.
C.
CODEN: 69DSA4

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Ionic liquids as green solvents for the dissolution
and regeneration of cellulose

AU Swatloski, Richard P.; Spear, Scott K.; Holbrey, John D.;
Rogers, Robin D.

AB With increasing governmental regulations restricting the use of current

cellulose solvents, the need to replace them is becoming more important. Ionic Liqs. (ILs) have gained considerable attention for their potential use as green solvents, and the use of ILs as -green' replacements. . . solvents has been studied in recent literature. In our labs., we have utilized ILs as solvents, for the dissoln. of cellulose. We have successfully dissolved, without pretreatment or derivitization, up to 30% wt/wt of cellulose in ILs, which enables the use of ILs as a feasible and effective non-volatile alternatives to some of the environmentally undesirable solvent systems currently in use. The cellulose can be regenerated from the ionic liquid by simply contacting them with water. This allows a simple, benign system for the processing of cellulose into fibers, monoliths and films by forming into an aqueous phase. Results from regenerated cellulose, as well as important intermol. forces responsible for the dissoln. of cellulose will be presented. This research is sponsored by The PG Research Foundation, Inc.

L9 ANSWER 24 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:617063 CAPLUS

TITLE: Ionic liquids: New solvents for nonderivitized cellulose dissolution

AUTHOR(S): Swatloski, Richard P.; Spear, Scott K.; Holbrey, John D.; Rogers, Robin D.

CORPORATE SOURCE: Department of Chemistry and Center for Green Manufacturing, The University of Alabama, Tuscaloosa, AL, 35487, USA

SOURCE: Abstracts of Papers, 224th ACS National Meeting, Boston, MA, United States, August 18-22, 2002 (2002), IEC-076. American Chemical Society: Washington, D. C. CODEN: 69CZPZ

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

TI Ionic liquids: New solvents for nonderivitized cellulose dissolution

AU Swatloski, Richard P.; Spear, Scott K.; Holbrey, John D.; Rogers, Robin D.

AB There are only a limited number of solvents that can effectively dissolve cellulose without derivitization; all have environmental downsides, and in some cases are even poor systems for the dissoln. Though it was first suggested in 1934 by Graenacher that molten N-ethylpyridinium chloride could be used to dissolve cellulose, at the time this seemed very impractical and of little value since the molten salt was, at the time, esoteric and a relatively high m.p. of 118 °C. Here we examine the history of cellulose solvents, the solubility of cellulose in ionic liqs. without activation or pretreatment, and the regeneration of cellulose from simple, nonvolatile ILs.

L9 ANSWER 25 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:287553 CAPLUS

DOCUMENT NUMBER: 136:387576

TITLE: Dissolution of cellulose with ionic liquids

AUTHOR(S): Swatloski, Richard P.; Spear, Scott K.; Holbrey, John D.; Rogers, Robin D.

CORPORATE SOURCE: Center for Green Manufacturing and Department of Chemistry, The University of Alabama, Tuscaloosa, AL,

SOURCE: 35487, USA
Journal of the American Chemical Society (2002),
124(18), 4974-4975
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Dissolution of cellulose with ionic liquids
AU Swatloski, Richard P.; Spear, Scott K.; Holbrey, John D.;
Rogers, Robin D.
AB Initial results that demonstrate that cellulose can be dissolved
without activation or pretreatment in, and regenerated from,
1-butyl-3-methylimidazolium chloride and other hydrophilic ionic
liqs. are reported. This may enable the application of
ionic liqs. as alternatives to environmentally
undesirable solvents currently used for dissoln. of this important
bio-resource.
ST butylmethylimidazolium chloride hydrophilic ionic liq
solvent effect cellulose dissoln
IT Cellulose pulp
Dissolution
Ionic liquids
Solvent effect
Thermal decomposition
(dissoln. of cellulose with hydrophilic ionic
liquid solvents)
IT 9004-34-6, Cellulose, processes
RL: PEP (Physical, engineering or chemical process); PYP (Physical
process); PROC (Process)
(dissoln. of cellulose with hydrophilic ionic
liquid solvents)
IT 64697-40-1 79917-90-1, 1-Butyl-3-methylimidazolium chloride
85100-77-2, 1-Butyl-3-methylimidazolium bromide 171058-17-6
174501-64-5 174501-65-6 344790-87-0, 1-Butyl-3-methylimidazolium
thiocyanate
RL: NUU (Other use, unclassified); USES (Uses)
(solvent; dissoln. of cellulose with hydrophilic
ionic liquid solvents)
L9 ANSWER 26 OF 26 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2001:639079 CAPLUS
TITLE: Derivatization of chitin in room temperature
ionic liquids
AUTHOR(S): Reichert, W. Matthew; Visser, Ann E.; Swatloski,
Richard P.; Spear, Scott K.; Rogers, Robin D.
CORPORATE SOURCE: Department of Chemistry and Center for Green
Manufacturing, The University of Alabama, Tuscaloosa,
AL, 35487, USA
SOURCE: Abstracts of Papers, 222nd ACS National Meeting,
Chicago, IL, United States, August 26-30, 2001 (2001),
IEC-025. American Chemical Society: Washington, D. C.
CODEN: 69BUZP
DOCUMENT TYPE: Conference; Meeting Abstract
LANGUAGE: English
TI Derivatization of chitin in room temperature ionic

liquids

- AU Reichert, W. Matthew; Visser, Ann E.; Swatloski, Richard P.;
Spear, Scott K.; Rogers, Robin D.
- AB . . . sepns. are usually the key steps preventing economic processing
of biomass. Chitin is the world's second most abundant biopolymer, behind
cellulose, and in this presentation, we will demonstrate the use
of room temperature ionic liqs. as solvents in the chemical
modification of chitin. In addition, the use of ionic liqs
. for value added processing (e.g., decolorization of chitin) will be
discussed.

=> file stng

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	91.70	120.10
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-19.50	-24.18

FILE 'STNGUIDE' ENTERED AT 16:01:40 ON 26 NOV 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Nov. 23, 2007 (20071123/UP).

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.18	120.28
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-24.18

FILE 'CAPLUS' ENTERED AT 16:03:16 ON 26 NOV 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 26 Nov 2007 VOL 147 ISS 23
FILE LAST UPDATED: 25 Nov 2007 (20071125/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.

They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> "ionic liquid" and cellulose

284921 "IONIC"
511 "IONICS"
285185 "IONIC"
("IONIC" OR "IONICS")
800777 "LIQUID"
138530 "LIQUIDS"
904338 "LIQUID"
("LIQUID" OR "LIQUIDS")
1105042 "LIQ"
104871 "LIQS"
1145172 "LIQ"
("LIQ" OR "LIQS")
1588658 "LIQUID"
("LIQUID" OR "LIQ")
11079 "IONIC LIQUID"
("IONIC"(W)"LIQUID")
360710 CELLULOSE
4428 CELLULOSES
361213 CELLULOSE
(CELLULOSE OR CELLULOSES)

L10 177 "IONIC LIQUID" AND CELLULOSE

=> 110 and ether?

603706 ETHER?

L11 11 L10 AND ETHER?

=> d 111 1-11 ibib kwic

L11 ANSWER 1 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:1088611 CAPLUS

DOCUMENT NUMBER: 147:387771

TITLE: Methods for modifying cellulosic polymers in
ionic liquids

INVENTOR(S): Scheibel, Jeffrey John; Kenneally, Corey James;
Menkhaus, Julie Ann; Seddon, Kenneth Richard; Chwala,
Prezemyslaw

PATENT ASSIGNEE(S): The Procter & Gamble Company, USA

SOURCE: U.S. Pat. Appl. Publ., 8pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
US 2007225190	A1	20070927	US 2007-726609	20070322
WO 2007112382	A1	20071004	WO 2007-US65000	20070327
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK,			

MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO,
 RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT,
 TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,
 GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
 BY, KG, KZ, MD, RU, TJ, TM

PRIORITY APPLN. INFO.:

US 2006-786415P

P 20060327

TI Methods for modifying cellulosic polymers in ionic liquids

AB Sulfation or sulfonation of cellulose and cellulose ethers is conducted in an ionic liquid such as a quaternary ammonium salt. Detergent compns. containing the sulfated or sulfonated reaction product are suitable for fabric cleansing.

ST cellulosic polymer modification ionic liq sulfonation sulfation

IT Sulfates, uses

RL: TEM (Technical or engineered material use); USES (Uses)

(alkyl ethoxy, detergents; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Sulfonic acids, uses

RL: TEM (Technical or engineered material use); USES (Uses)

(arenesulfonic, salts, detergents; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Catalysts

(bleach; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Zeolites (synthetic), uses

RL: TEM (Technical or engineered material use); USES (Uses)

(builders; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Alcohols, uses

RL: TEM (Technical or engineered material use); USES (Uses)

(ethoxylated, detergents; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Detergents

(laundry; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Antibacterial agents

Bleaching agents

Brightening agents

Chelating agents

Creaseproofing

Detergent builders

Dyes

Fireproofing agents

Ionic liquids

Perfumes

Sizing

Skin conditioners

Soils

Sulfation

Sulfonation

Surfactants
 Textiles
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Enzymes, uses
 RL: CAT (Catalyst use); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Clays, processes
 RL: REM (Removal or disposal); PROC (Process)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Vitamins
 RL: SPN (Synthetic preparation); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Biopolymers
 Quaternary ammonium compounds, uses
 RL: TEM (Technical or engineered material use); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT UV radiation
 (protection; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT Aromatic compounds
 RL: TEM (Technical or engineered material use); USES (Uses)
 (sulfonates, detergents; methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT 9004-34-6, Cellulose, uses
 RL: PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PROC (Process); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT 9032-43-3P, Cellulose sulfate
 RL: SPN (Synthetic preparation); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

IT 124-41-4, Sodium methoxide 5329-14-6, Sulfamic acid 7664-93-9,
 Sulfuric acid, uses 7790-94-5, Chlorosulfonic acid 28322-92-1, Sulfur
 trioxide-pyridine complex 79917-90-1, 1-n-Butyl-3-methylimidazolium
 chloride
 RL: TEM (Technical or engineered material use); USES (Uses)
 (methods for modifying cellulosic polymers in ionic liqs. and their applications for fabric cleansing)

L11 ANSWER 2 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:933138 CAPLUS

DOCUMENT NUMBER: 147:290978

TITLE: Method of processing a biological and/or chemical sample

INVENTOR(S): Pipper, Juergen; Hsieh, Tseng-Ming; Neuzil, Pavel

PATENT ASSIGNEE(S): Agency for Science, Technology and Research, Singapore

SOURCE: PCT Int. Appl., 67pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2007094739	A1	20070823	WO 2006-SG29	20060213
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.: WO 2006-SG29 20060213
 REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

IT Air analysis
 Amniotic fluid
 Animal tissue
 Animal tissue culture
 Bioluminescence
 Blood analysis
 Blood plasma
 Blood serum
 Body fluid
 Bone marrow
 Cell
 Ceramics
 Cerebrospinal fluid
 Colorimetry
 Composites
 Cytolysis
 Drops
 Dyes
 Electromagnetic field
 Electromagnets
 Environmental analysis
 Enzyme-linked immunosorbent assay
 Extracellular matrix
 Extraction
 Feces
 Films
 Filtration
 Fluorometry
 Food analysis
 Groundwaters
 Hair
 Immobilization, molecular or cellular
 Interferometry
Ionic liquids

Ions
Leukocyte
Lymph
Magnetic field
Magnetic fluids
Magnets
Microorganism
Milk analysis
Mixing
Nail (anatomical)
Neoplasm
Optical diffraction
Paper
Particles
Pharmaceutical analysis
Photometry
Radioactive fallout
Rainwater
Sample preparation
Semen
Skin
Soil analysis
Spectroscopy
Sputum
Urine analysis
Virus
Wastewater

(method of processing biol. and/or chemical sample in fluid droplet)

IT Agglutinins and Lectins
Aluminosilicates, uses
Ankyrins
Antibodies and Immunoglobulins
Calmodulins
Crown ethers
Enzymes, uses
Gelatins, uses
Ligands
Lipocalins
Nucleic acids
Peptides, uses
Protein A
Proteins

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(method of processing biol. and/or chemical sample in fluid droplet)

IT Ethers, analysis

RL: ARU (Analytical role, unclassified); ANST (Analytical study)
(method of processing biol. and/or chemical sample in fluid droplet)

IT 53-59-8, NADP 53-84-9, NAD 56-87-1, Lysine, uses 69-79-4, Maltose
70-18-8, Glutathione, uses 74-79-3, Arginine, uses 618-39-3,
Benzamidine 1398-61-4, Chitin 1406-11-7, Polymyxin 9004-34-6,
Cellulose, uses 9005-49-6, Heparin, uses

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(method of processing biol. and/or chemical sample in fluid droplet)

L11 ANSWER 3 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2007:458864 CAPLUS
DOCUMENT NUMBER: 146:458065

TITLE: The application using non-covalent bond between a cucurbituril derivative and a ligand

INVENTOR(S): Kim, Kimoon; Baek, Kangkyun; Kim, Jeeyoun; Hwang, Ilha; Ko, Young-Ho; Selvapalam, Narayanan; Nagarajan, Erumaipatty R.; Park, Kyeng-Min

PATENT ASSIGNEE(S): Postech Academy-Industry Foundation, S. Korea

SOURCE: PCT Int. Appl., 67pp.
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2007046575	A1	20070426	WO 2006-KR687	20060228
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
KR 2007050747	A	20070516	KR 2006-18434	20060224
US 2007092867	A1	20070426	US 2006-407143	20060420
PRIORITY APPLN. INFO.:				
			KR 2005-99379	A 20051020
			KR 2005-108312	A 20051112
			KR 2006-891	A 20060104
			KR 2006-18434	A 20060224

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB . . . phase, a biomol., an antioxidant, a chemical therapeutic agent, an antihistaminic agent, a cucurbituril dendrimer, a cyclodextrin derivative, a crown ether derivative, a calixarene derivative, a cyclophane derivative, a cyclic peptide derivative, a metallic ion, a chromophore, a fluorescent material, a . . .

IT Affinity chromatographic stationary phases

Affinity chromatography

Amino group

Antihistamines

Antioxidants

Antitumor agents

Catalysts

Cations

Cell

Cell death

Cell membrane

Chemical formula

Chloromethylation

Chromophores

Fluorescent substances

Immobilization, molecular or cellular

Ionic liquids

Magnetic materials

Mixtures

Nanotubes

Nanowires

Noncovalent bond

Phosphors

Radioactive substances

Separation

Solvents

Test kits

Virus

(application using non-covalent bond between cucurbituril derivative and ligand)

IT Agglutinins and Lectins

Amino acids, uses

Antibodies and Immunoglobulins

Antigens

Biochemical compounds

Chemical compounds

Coenzymes

Crown ethers

Cyclic peptides

Cyclophanes

Enzymes, uses

Fatty acids, uses

Glass, uses

Glycoproteins

Histones

Hormones, animal, uses

Ligands

Metallocenes

Metals, uses

Nucleic acids

Polymers, uses

Polysaccharides, uses

Receptors

Resins

Vitamins

RL: NUU (Other use, unclassified); USES (Uses)

(application using non-covalent bond between cucurbituril derivative and ligand)

IT 52-90-4, L-Cysteine, uses 56-65-5, 5'-ATP, uses 58-64-0, 5'-ADP, uses
60-29-7, Diethyl ether, uses 64-17-5, Ethanol, uses 67-56-1,
Methanol, uses 67-66-3, Chloroform, uses 67-68-5, Dimethyl sulfoxide,
uses 68-12-2, Dimethylformamide, uses 70-18-8, Glutathione, uses
71-00-1, L-Histidine, uses 73-22-3, L-Tryptophan, uses 75-05-8,
Acetonitrile, uses 75-09-2, Methylene chloride, uses 76-05-1,
Trifluoroacetic acid, uses 108-88-3, Toluene, uses 108-90-7,
Chlorobenzene, uses 109-99-9, Tetrahydrofuran, uses 110-86-1,
Pyridine, uses 111-46-6, Diglycol, uses 120-94-5, N-Methylpyrrolidine
121-44-8, Triethylamine, uses 123-91-1, Dioxane, uses 124-38-9, Carbon
dioxide, uses 281-23-2, Adamantane 768-94-5, Adamantanamine
1314-23-4, Zirconium oxide, uses 1330-20-7, Xylene, uses 1336-21-6,
Ammonium hydroxide 7440-21-3, Silicon, uses 7732-18-5, Water, uses
9000-92-4, Amylase 9001-54-1, Hyaluronidase 9001-92-7, Proteinase
9002-10-2, Phenoloxidase 9003-99-0, Peroxidase 9004-34-6,

Cellulose, uses 9012-36-6, Sepharose 9012-54-8, Cellulase 9013-79-0, Esterase 9025-56-3, Hemicellulase 9029-60-1, Lipxygenase 9032-75-1, Pectinase 9035-73-8, Oxidase 9037-17-6 9037-80-3, Reductase 9067-74-7, Arabinosidase 9075-68-7, Pullulanase 12176-38-4, Ferrocene methylamine 12619-70-4D, Cyclodextrin, derivs. 37278-89-0, Xylanase 37341-53-0, Keratinase 42613-30-9, Ligninase 51377-41-4, Cutinase 54724-00-4, Curdlan 80262-44-8D, Cucurbituril, derivs.

RL: NUU (Other use, unclassified); USES (Uses)

(application using non-covalent bond between cucurbituril derivative and ligand)

L11 ANSWER 4 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1262324 CAPLUS

DOCUMENT NUMBER: 146:483399

TITLE: Preparation method of cellulose ether derivative in 1-alkyl-3-alkyl-imidazolinium ionic liquid

INVENTOR(S): Park, Young Seok; Park, Jung Ho

PATENT ASSIGNEE(S): Kolon Industries, Inc., S. Korea

SOURCE: Repub. Korean Kongkae Taeho Kongbo, No pp. given
CODEN: KRXXA7

DOCUMENT TYPE: Patent

LANGUAGE: Korean

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
KR 2006086069	A	20060731	KR 2005-6937	20050126
PRIORITY APPLN. INFO.:			KR 2005-6937	20050126

TI Preparation method of cellulose ether derivative in 1-alkyl-3-alkyl-imidazolinium ionic liquid

AB Provided is a method for preparing a cellulose ether derivative under the homogeneous reaction condition to minimize the amount of remaining reaction solvent contained in the final product. The method comprises the steps of dissolving cellulose in an imidazolinium-based ionic compound such as a 1-alkyl-3-alkyl-imidazolinium salt; and etherifying using a metal hydroxide as a catalyst to prepare a cellulose ether derivative. Preferably the alkali cellulose activated by the addition of the metal hydroxide catalyst is prepared into cellulose ether by adding an etherification agent. Preferably the metal hydroxide is at least one selected from LiOH, NaOH, KOH, Ca(OH)₂, Mg(OH)₂, Al(OH)₃ and nickel hydroxide.

ST cellulose ether manuf etherification
imidazolinium ionic liq

IT Hydroxides (inorganic)

RL: CAT (Catalyst use); USES (Uses)

(catalyst; preparation method of cellulose ether derivative in 1-alkyl-3-alkyl-imidazolinium ionic liquid)

IT Ionic liquids

(preparation method of cellulose ether derivative in 1-alkyl-3-alkyl-imidazolinium ionic liquid)

IT 1305-62-0, Calcium hydroxide, uses 1309-42-8, Magnesium hydroxide 1310-58-3, Potassium hydroxide, uses 1310-65-2, Lithium hydroxide 1310-73-2, Sodium hydroxide, uses 12054-48-7, Nickel hydroxide

21645-51-2, Aluminum hydroxide, uses
 RL: CAT (Catalyst use); USES (Uses)
 (catalyst; preparation method of cellulose ether derivative
 in 1-alkyl-3-alkyl-imidazolinium ionic liquid)

IT 9004-34-6DP, Cellulose, ether
 RL: IMF (Industrial manufacture); PREP (Preparation)
 (preparation method of cellulose ether derivative in
 1-alkyl-3-alkyl-imidazolinium ionic liquid)

IT 288-32-4D, Imidazole, N-alkyl salts
 RL: NUU (Other use, unclassified); USES (Uses)
 (preparation method of cellulose ether derivative in
 1-alkyl-3-alkyl-imidazolinium ionic liquid)

L11 ANSWER 5 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:367288 CAPLUS
 DOCUMENT NUMBER: 144:398457
 TITLE: Indicator device having an active agent encapsulated
 in an electrospun nanofiber
 INVENTOR(S): McDonnell, Gerald E.; Fiorello, Anthony; Smith, Daniel
 J.
 PATENT ASSIGNEE(S): Steris Inc., USA
 SOURCE: U.S. Pat. Appl. Publ., 10 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2006083657	A1	20060420	US 2004-965350	20041014
AU 2005333237	A1	20061228	AU 2005-333237	20050909
CA 2582979	A1	20061228	CA 2005-2582979	20050909
WO 2006137848	A2	20061228	WO 2005-US32448	20050909
WO 2006137848	A3	20070802		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
EP 1799805	A2	20070627	EP 2005-858110	20050909
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU				
IN 2007MN00501	A	20070803	IN 2007-MN501	20070405
PRIORITY APPLN. INFO.:			US 2004-965350	A 20041014
			WO 2005-US32448	W 20050909
IT Antimicrobial agents				
Biocides				
Cations				

Chemiluminescence spectroscopy

Cyanine dyes

Fluorometry

Indicators

Ionic liquids

Microelectrodes

Mycobacterium

Nanofibers

Phosphorescence

Plasticizers

Waters

(indicator device having active agent encapsulated in electrospun nanofiber)

IT Azo compounds

Carbonyl complexes

Carotenes, biological studies

Crown ethers

Nitro compounds

Nitroso compounds

Prion proteins

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(indicator device having active agent encapsulated in electrospun nanofiber)

IT 9002-89-5, Polyvinyl alcohol 9002-98-6 9003-05-8, Polyacrylamide

9003-39-8 9004-34-6, Cellulose, biological studies

9004-57-3, Ethylcellulose 24980-41-4, Polycaprolactone 25248-42-4, Polycaprolactone 26913-06-4, Poly[imino(1,2-ethanediyl)] 76600-67-4, Tecoflex

RL: BUU (Biological use, unclassified); DEV (Device component use); BIOL (Biological study); USES (Uses)

(indicator device having active agent encapsulated in electrospun nanofiber)

L11 ANSWER 6 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:1255766 CAPLUS

DOCUMENT NUMBER: 143:488016

TITLE: Ionic liquids used as solvents in titrimetric analysis

INVENTOR(S): Bosmann, Andreas; Schubert, Thomas Juergen Siegfried

PATENT ASSIGNEE(S): Germany

SOURCE: Brit. UK Pat. Appl., 17 pp.

CODEN: BAXXDU

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
GB 2414553	A	20051130	GB 2005-10772	20050526
DE 102004025756	A1	20051215	DE 2004-102004025756	20040526
US 2005287677	A1	20051229	US 2005-136900	20050525
JP 2005338092	A	20051208	JP 2005-154529	20050526
PRIORITY APPLN. INFO.:			DE 2004-102004025756A	20040526
REFERENCE COUNT:	2	THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT		

TI Ionic liquids used as solvents in titrimetric analysis

AB Ionic liqs. provide new solvents for the titrimetric anal. of substances or mixts. of substances which are insol. or poorly soluble in conventional solvents, such as e.g. proteins, cellulose, etc. This allows direct homogeneous titration of ingredients of these substances. In addition to the substances which are to be analyzed the titration reagents may advantageously also be dissolved in an ionic liquid

ST ionic liq solvent titrimetry

IT Named reagents and solutions
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses) (Karl Fischer's; ionic liqs. as solvents for titrimetric anal.)

IT Cheese
 (Parmesan; ionic liqs. as solvents for titrimetric anal.)

IT Sulfoxides
 RL: ARU (Analytical role, unclassified); ANST (Analytical study) (alkyl, dialkylsulfoxides; ionic liqs. as solvents for titrimetric anal.)

IT Fats and Glyceridic oils, analysis
 RL: AMX (Analytical matrix); ANST (Analytical study) (butter; ionic liqs. as solvents for titrimetric anal.)

IT Titration
 (conductometric; ionic liqs. as solvents for titrimetric anal.)

IT Titration
 (coulometric; ionic liqs. as solvents for titrimetric anal.)

IT Alkanes, analysis
 RL: ARU (Analytical role, unclassified); ANST (Analytical study) (halo; ionic liqs. as solvents for titrimetric anal.)

IT Aromatic compounds
 RL: ARU (Analytical role, unclassified); ANST (Analytical study) (haloaroms.; ionic liqs. as solvents for titrimetric anal.)

IT Corylus avellana
 Food analysis
 Hygroscopicity
 Indicators
Ionic liquids
 NMR spectroscopy
 Solvents
 Titration
 (ionic liqs. as solvents for titrimetric anal.)

IT Alcohols, analysis
 Alkanes, analysis
 Amides, analysis
 Amines, analysis
 Aromatic compounds
Ethers, analysis
 Phosphonium compounds
 Quaternary ammonium compounds, analysis
 RL: ARU (Analytical role, unclassified); ANST (Analytical study) (ionic liqs. as solvents for titrimetric anal.)

IT Titration
(potentiometric; ionic liqs. as solvents for
titrimetric anal.)

IT Titration
(spectrophotometric; ionic liqs. as solvents for
titrimetric anal.)

IT Titration
(thermometric; ionic liqs. as solvents for
titrimetric anal.)

IT 7732-18-5, Water, analysis
RL: ANT (Analyte); ANST (Analytical study)
(ionic liqs. as solvents for titrimetric anal.)

IT 174899-82-2 258864-54-9, Trihexyltetradecylphosphonium chloride
342573-75-5 792188-85-3
RL: ARU (Analytical role, unclassified); ANST (Analytical study)
(ionic liqs. as solvents for titrimetric anal.)

L11 ANSWER 7 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:523500 CAPLUS

DOCUMENT NUMBER: 143:28326

TITLE: Etherification of cellulose in
ionic liquid solutions

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 23 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005054298	A1	20050616	WO 2004-FI730	20041202
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
FI 2003001763	A	20050604	FI 2003-1763	20031203
FI 116140	B1	20050930		
CA 2548007	A1	20050616	CA 2004-2548007	20041202
EP 1689788	A1	20060816	EP 2004-801227	20041202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS			
US 2007112185	A1	20070517	US 2007-581491	20070116
PRIORITY APPLN. INFO.:			FI 2003-1763	A 20031203
			WO 2004-FI730	W 20041202

OTHER SOURCE(S): MARPAT 143:28326

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- TI Etherification of cellulose in ionic liquid solutions
- AB Cellulose is mixed and dissolved in an ionic liquid solvent and the solution is treated with an etherifying agent in the presence of inorg. base to form a cellulose ether, which is subsequently separated from the solution. The dissoln. and the etherification are carried out in the absence of organic base and in the substantial absence of H₂O. Microwave irradiation and/or pressure can be applied to assist in dissoln. and etherification. Thus, 50 mg cellulose was dissolved in 5 g 1-butyl-3-methylimidazolium chloride (m. 60°) with the aid of microwaves to give 1% solution. ClCH₂CO₂H (2.05. . . equiv of solid NaOH, the reaction mixture was heated for 2 h at 100° under microwave radiation and the resulting CM-cellulose was precipitated with MeOH, washed with MeOH and 80% aqueous MeOH, and dried.
- ST cellulose etherification ionic liq solvent microwave; butylmethylimidazolium chloride solvent CM cellulose manuf; chloroacetic acid etherification cellulose butylmethylimidazolium chloride solvent
- IT Etherification
Ionic liquids
 (etherification of cellulose in ionic liquid solution)
- IT Microwave
 (etherification of cellulose in ionic liquid solution in presence of)
- IT 9004-32-4P, CM cellulose sodium salt
 RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
 (etherification of cellulose in ionic liquid solution)
- IT 9004-34-6, Cellulose, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (etherification of cellulose in ionic liquid solution)
- IT 79-11-8, Chloroacetic acid, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (etherification of cellulose;
etherification of cellulose in ionic liquid solution)
- IT 79917-90-1, 1-Butyl-3-methylimidazolium chloride
 RL: TEM (Technical or engineered material use); USES (Uses)
 (solvent; etherification of cellulose in ionic liquid solution)

L11 ANSWER 8 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:443826 CAPLUS

DOCUMENT NUMBER: 143:135076

TITLE: Ionic Liquids and Paper

AUTHOR(S): Przybysz, Kazimierz; Drzewinska, Ewa; Stanislawska, Anna; Wysocka-Robak, Agnieszka; Cieniecka-Roslonkiewicz, Anna; Foksowicz-Flaczyk, Joanna; Pernak, Juliusz

CORPORATE SOURCE: Institute of Papermaking and Printing, Lodz University of Technology, Lodz, 90-924, Pol.

SOURCE: Industrial & Engineering Chemistry Research (2005), 44(13), 4599-4604

CODEN: IECRED; ISSN: 0888-5885
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- TI Ionic Liquids and Paper
- AB The influence of ionic liqs. (ILs) on the characteristics of paper was studied. The ILs used include 3-alkyl-1-methylimidazolium tetrafluoroborates and prepared 3-alkoxymethyl-1-methylimidazolium tetrafluoroborates and 3-alkoxymethyl-1-methylimidazolium. . . impregnate paper, affecting its strength and other phys. parameters. The IL-treated paper had decreased strength, which resulted from weakening of cellulose hydrogen bonds and the paper wettability improved. Paper treated with 1-methyl-3-octyloxymethylimidazolium tetrafluoroborate proved to be fully resistant to activity of. . .
- ST paper treatment ionic liq methylimidazolium tetrafluoroborate trifluoromethanesulfonylimide strength; mold resistant paper ionic liq treatment
- IT Alternaria alternata
 Aspergillus amstelodami
 Aspergillus niger
 Aspergillus terreus
 Aureobasidium pullulans
 Chaetomium globosum
 Cladosporium herbarum
 Fungicides
Ionic liquids
 Paecilomyces variotii
 Paper
 Penicillium brevi-compactum
 Penicillium funiculosum
 Penicillium ochro-chloron
 Scopulariopsis brevicaulis
 Stachybotrys chartarum
 Trichoderma viride
 Wettability
 (mech. properties and resistant to mold and fungus of paper treated with methylimidazolium based ionic liqs.)
- IT 99874-27-8P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; mech. properties and resistant to mold and fungus of paper treated with methylimidazolium based ionic liqs .)
- IT 616-47-7, 1-Methylimidazole 39979-92-5, Chloromethylhexyl ether
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (mech. properties and resistant to mold and fungus of paper treated with methylimidazolium based ionic liqs.)
- IT 244193-52-0P 244193-56-4P 350701-79-0P 350701-81-4P 350701-83-6P
 350701-85-8P 350701-87-0P 852951-58-7P 859213-79-9P 859213-80-2P
 859213-81-3P 859213-82-4P 859213-83-5P
 RL: SPN (Synthetic preparation); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)
 (mech. properties and resistant to mold and fungus of paper treated with methylimidazolium based ionic liqs.)

ACCESSION NUMBER: 2005:270816 CAPLUS

DOCUMENT NUMBER: 143:479529

TITLE: Unconventional dissolution and derivatization of cellulose

AUTHOR(S): Fischer, Steffen

CORPORATE SOURCE: Fraunhofer Institute of Applied Polymer Research, Potsdam, 14476, Germany

SOURCE: Lenzinger Berichte (2004), 83, 71-78

CODEN: LEBEAW; ISSN: 0024-0907

PUBLISHER: Lenzing AG

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Unconventional dissolution and derivatization of cellulose

AB The preparation of cellulose solns. is important for derivatization and blend formation of the natural polymer. Besides solvents like CS₂/NaOH and NMNO*H₂O unconventional solvent systems can be applied for dissoln. of cellulose. This group of solvents includes inorg. molten salts and ionic liqs. Inorg. molten salts can be used as efficient solvents for cellulose in a wide range of d.p. Furthermore molten salts can be applied as reaction medium for the derivatization of cellulose. For both dissoln. and derivatization of cellulose the knowledge of the solution state as well as information about chemical interactions with the solvent system is essential. Using the melts of LiClO₄·3H₂O, NaSCN/KSCN/LiSCN·2H₂O and LiCl/ZnCl₂/H₂O as cellulose solvents factors which determine the dissolving ability will be discussed. Besides the specific structure of the molten salt hydrate, the . . . for the dissolving capability of a molten salt hydrate system. The application of inorg. molten salts as a medium for cellulose functionalization is demonstrated for cellulose carboxymethylation and acetylation.

ST cellulose dissoln derivatization inorg molten salt; CM cellulose prepn inorg molten salt medium; acetylcellulose prepn inorg molten salt medium

IT Dissolution

Esterification

Etherification

(unconventional dissoln. and derivatization of cellulose using inorg. molten salts)

IT 333-20-0, Potassium thiocyanate 540-72-7, Sodium thiocyanate 7447-41-8, Lithium chloride, uses 7646-85-7, Zinc chloride, uses 13453-78-6, Lithium perchlorate trihydrate 84372-58-7, Lithium thiocyanate dihydrate

RL: NUU (Other use, unclassified); USES (Uses)

(unconventional dissoln. and derivatization of cellulose using inorg. molten salts)

IT 9004-34-6, Cellulose, processes

RL: PEP (Physical, engineering or chemical process); PYP (Physical process); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)

(unconventional dissoln. and derivatization of cellulose using inorg. molten salts)

IT 9004-32-4P, CM cellulose sodium salt 9004-35-7P, Acetylcellulose

RL: SPN (Synthetic preparation); PREP (Preparation)
(unconventional dissoln. and derivatization of cellulose
using inorg. molten salts)

L11 ANSWER 10 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:158715 CAPLUS
DOCUMENT NUMBER: 142:242565
TITLE: Dissolution and delignification of lignocellulosic
materials with ionic liquid
solvent under microwave irradiation
INVENTOR(S): Myllymäeki, Vesa; Aksela, Reijo
PATENT ASSIGNEE(S): Kemira Oyj, Finland
SOURCE: PCT Int. Appl., 25 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE			
-----	----	-----	-----	-----			
WO 2005017001	A1	20050224	WO 2004-FI476	20040813			
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW						
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG						
FI 2003001156	A	20050216	FI 2003-1156	20030815			
FI 115835	B1	20050729					
CA 2532989	A1	20050224	CA 2004-2532989	20040813			
EP 1654307	A1	20060510	EP 2004-742219	20040813			
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK						
BR 2004013435	A	20061010	BR 2004-13435	20040813			
PRIORITY APPLN. INFO.:			FI 2003-1156	A 20030815			
			WO 2004-FI476	W 20040813			
OTHER SOURCE(S):	MARPAT 142:242565						
REFERENCE COUNT:	4	THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT					
TI	Dissolution and delignification of lignocellulosic materials with <u>ionic liquid</u> solvent under microwave irradiation						
AB	Wood, straw, and other natural lignocellulosic materials can be dissolved in an <u>ionic liquid</u> solvent under microwave irradiation and/or under pressure, and <u>cellulose</u> and other organic compds., such as lignin and extractives, can also be separated from the solution by precipitating with non-solvent, such as water, alcs., ketones, and <u>ethers</u> , of <u>cellulose</u> . Thus, plywood sawdust was dissolved in 1-butyl-3-methyl-imidazolium chloride under microwave irradiation						
ST	dissoln delignification lignocellulosic <u>ionic liq</u> solvent microwave irradsn; plywood sawdust wood straw						

butylmethylimidazolium chloride dissoln microwave irradsn

IT Wood
 (chips; dissoln. and delignification of lignocellulosic materials with
ionic liquid solvent under microwave irradiation)

IT Dissolution
 Straw
 Wood
 (dissoln. and delignification of lignocellulosic materials with
ionic liquid solvent under microwave irradiation)

IT Solvents
 (ionic, liquid; dissoln. and delignification of
 lignocellulosic materials with ionic liquid solvent
 under microwave irradiation)

IT Microwave
 (irradiation; dissoln. and delignification of lignocellulosic materials
 with ionic liquid solvent under microwave irradiation)

IT Wood boards
 (plywood, sawdust; dissoln. and delignification of lignocellulosic
 materials with ionic liquid solvent under microwave
 irradiation)

IT Sawdust
 (plywood; dissoln. and delignification of lignocellulosic materials
 with ionic liquid solvent under microwave irradiation)

IT Wood
 (soft; dissoln. and delignification of lignocellulosic materials with
ionic liquid solvent under microwave irradiation)

IT Alcohols, uses
Ethers, uses
 Ketones, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (solvent; dissoln. and delignification of lignocellulosic materials
 with ionic liquid solvent under microwave irradiation)

IT 9004-34-6P, Cellulose, preparation 9005-53-2P, Lignin,
 preparation
 RL: PUR (Purification or recovery); PREP (Preparation)
 (dissoln. and delignification of lignocellulosic materials with
ionic liquid solvent under microwave irradiation)

IT 79917-90-1, 1-Butyl-3-methyl-imidazolium chloride
 RL: NUU (Other use, unclassified); USES (Uses)
 (solvent; dissoln. and delignification of lignocellulosic materials
 with ionic liquid solvent under microwave irradiation)

L11 ANSWER 11 OF 11 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:802385 CAPLUS

DOCUMENT NUMBER: 141:298755

TITLE: Ionically conductive membranes for protection of
 active metal anodes and battery cells

INVENTOR(S): Visco, Steven J.; Nimon, Yevgeniy S.; Katz, Bruce D.

PATENT ASSIGNEE(S): Polyplus Battery Company, USA

SOURCE: U.S. Pat. Appl. Publ., 25 pp., Cont.-in-part of U.S.
 Ser. No. 731,771.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004191617	A1	20040930	US 2004-772228	20040203
US 2004126653	A1	20040701	US 2003-686189	20031014
US 7282296	B2	20071016		
US 2004142244	A1	20040722	US 2003-731771	20031205
US 7282302	B2	20071016		
WO 2005038962	A2	20050428	WO 2004-US33372	20041008
WO 2005038962	A3	20051229		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2005100793	A1	20050512	US 2004-986441	20041110
PRIORITY APPLN. INFO.:			US 2002-418899P	P 20021015
			US 2003-511710P	P 20031014
			US 2003-686189	A2 20031014
			US 2003-518948P	P 20031110
			US 2003-731771	A2 20031205
			US 2004-772228	A 20040203
IT	Battery anodes Ceramics Gelation agents Glass ceramics Ionic liquids Primary batteries Secondary batteries (ionically conductive membranes for protection of active metal anodes and battery cells)			
IT	Esters, uses Ethers , uses Fluoropolymers, uses Halides Metallic glasses Nitrides Phosphonium compounds Polyoxyalkylenes, uses Polysulfides RL: DEV (Device component use); USES (Uses) (ionically conductive membranes for protection of active metal anodes and battery cells)			
IT	79-20-9, Methyl acetate 96-47-9, 2-Methyltetrahydrofuran 105-58-8, Diethyl carbonate 107-31-3, Methyl formate 109-99-9, Thf, uses 110-71-4, 1,2-Dimethoxyethane 463-79-6D, Carbonic acid, organic esters 616-38-6, Dimethyl carbonate 623-53-0, Ethyl methyl carbonate 646-06-0, 1,3-Dioxolane 1072-47-5, 1,3-Dioxolane, 4-methyl- 1313-13-9, Manganese dioxide, uses 1313-27-5, Molybdenumoxide moo3, uses 1314-62-1, Vanadium oxide (V2O5), uses 1317-37-9, Iron sulfide Fes 1317-38-0, Copper oxide (CuO), uses 1317-40-4, Copper sulfide Cus , 7439-93-2, Lithium, uses 7439-93-2D, Lithium, intercalation compound			

7447-41-8, Lithium chloride (LiCl), uses 7550-35-8, Lithium bromide (LiBr) 7704-34-9, Sulfur, uses 7784-01-2, Silver chromate 7789-24-4, Lithium fluoride, uses 9004-67-5, Methyl cellulose 10377-51-2, Lithium iodide 11105-02-5, Silver vanadium oxide 12037-42-2, Vanadium oxide v6o13 12039-13-3, Titanium sulfide (TiS2) 12057-29-3, Lithium phosphide li3p 12068-85-8, Iron sulfide fes2 12789-09-2, Copper vanadium oxide 15365-14-7, Iron lithium phosphate felipo4 16969-45-2D, Pyridinium, derivs. 17009-90-4D, Imidazolium, derivs. 24937-79-9, PvdF 25014-41-9, Polyacrylonitrile 25322-68-3, Peo 26134-62-3, Lithium nitride (Li3N) 39300-70-4, Lithium nickeloxide 39457-42-6, Lithium manganese oxide 52627-24-4, Cobalt lithium oxide 70780-99-3, Lisicon 77641-62-4, Nasicon 155371-19-0, 1-Ethyl-3-methylimidazolium hexafluorophosphate 184905-46-2, Lithium nitrogen phosphorus oxide 244193-50-8, 1-Hexyl-3-methylimidazolium tetrafluoroborate 328090-25-1 445473-58-5, 1-Butyl-3-methylimidazolium octyl sulfate

RL: DEV (Device component use); USES (Uses)
 (ionically conductive membranes for protection of active metal anodes and battery cells)

=> d his

(FILE 'HOME' ENTERED AT 15:58:30 ON 26 NOV 2007)

FILE 'CAPLUS' ENTERED AT 15:58:43 ON 26 NOV 2007

E MYLLYMAKI/AU

L1 19 S E2

E AKSELA/AU

L2 47 S E10 OR E11

L3 60 DUP REMOVE L1 L2 (6 DUPLICATES REMOVED)

L4 19 S L3

L5 41 S L3

L6 6 S L3 AND IONIC

FILE 'STNGUIDE' ENTERED AT 16:00:24 ON 26 NOV 2007

FILE 'CAPLUS' ENTERED AT 16:00:34 ON 26 NOV 2007

E SWATLOSKI/AU

L7 91 S E4-E7

L8 83 S L7 AND "IONIC LIQUID"

L9 26 S L8 AND CELLULOSE

FILE 'STNGUIDE' ENTERED AT 16:01:40 ON 26 NOV 2007

FILE 'CAPLUS' ENTERED AT 16:03:16 ON 26 NOV 2007

L10 177 "IONIC LIQUID" AND CELLULOSE

L11 11 L10 AND ETHER?

=> l10 and microwave

123811 MICROWAVE

10771 MICROWAVES

125649 MICROWAVE

(MICROWAVE OR MICROWAVES)

L12 6 L10 AND MICROWAVE

=> d l12 1-6 ibib abs

L12 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:763873 CAPLUS
DOCUMENT NUMBER: 147:141573
TITLE: Processes for biomass treatment with higher energy efficiency
INVENTOR(S): Gurin, Michael H.
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 21pp., Cont.-in-part of U.S. Ser. No. 309,025.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2007161095	A1	20070712	US 2007-691070	20070326
WO 2007112090	A2	20071004	WO 2007-US7430	20070326
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

PRIORITY APPLN. INFO.:
US 2005-593485P P 20050118
US 2005-595167P P 20050613
US 2006-306911 A2 20060116
US 2006-767403P P 20060325
US 2006-309025 A2 20060612

AB A high efficiency method for synthesizing biomass fuels leveraging the synergistic impact of ionic liqs. on both the significant gains in pretreatment of biomass and the utilization of the combination of ionic liqs. and carbon dioxide under supercrit. conditions for energy generation is provided. The strategic use of heat exchangers, preferably microchannel heat exchangers and microchannel reactors further increase the efficiency and performance of the system by extensive heat recovery and the direct utilization of the biomass solution as the working fluid of a thermodyn. cycle.

L12 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:38602 CAPLUS
DOCUMENT NUMBER: 146:123422
TITLE: Ionic liquid reconstituted cellulose composites as solid support matrices with good transparency for biocatalytic reaction
INVENTOR(S): Rogers, Robin D.; Daly, Daniel T.; Turner, Megan B.; Spear, Scott K.; Holbrey, John D.
PATENT ASSIGNEE(S): The University of Alabama, USA
SOURCE: PCT Int. Appl., 73pp.

CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2007005388	A2	20070111	WO 2006-US24863	20060627
WO 2007005388	A3	20070329		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

US 2007006774	A1	20070111	US 2006-475630	20060627
PRIORITY APPLN. INFO.:			US 2005-694902P	P 20050629
OTHER SOURCE(S):	MARPAT 146:123422			

AB Disclosed are composites comprising regenerated cellulose, a first active substance, a second active substance, and a linker. Thus, microcryst. cellulose was dissolved in 1-butyl-3-methylimidazolium chloride using microwave pulse heating at 120-150°, cooled to 60° to form a super-cooled liquid, 20% (based on cellulose) poly(L-lysine hydrobromide) was added therein, homogenized, cast onto a glass plate, the resulting film soaked in water for at least 24 h to leach residual from the film to give a reconstituted cellulose film, showing good transparency.

L12 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:754555 CAPLUS

DOCUMENT NUMBER: 145:194651

TITLE: Method for complete enzymatic hydrolysis of straw cellulose pretreated with steam and microwave

INVENTOR(S): Chen, Hongzhang; Liu, Liying

PATENT ASSIGNEE(S): Institute of Process Engineering, Chinese Academy of Sciences, Peop. Rep. China

SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 7 pp.
CODEN: CNXXEV

DOCUMENT TYPE: Patent

LANGUAGE: Chinese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1806945	A	20060726	CN 2005-10011217	20050120
PRIORITY APPLN. INFO.:			CN 2005-10011217	20050120

AB The title method comprises: (1) steam-blasting straws with water content of 10-35% under steam pressure of 1.0-1.5 MPa for 2-7 min, (2) washing

with water of 50-100°C, drying, mixing with ionic liquid at a solid-liquid ratio of 1 : (5-50), and heating directly or by microwave under stirring for 5-60 min, (3) washing the treated straw with water, and (4) hydrolyzing with cellulase at below 50°C and pH 4.8 for 48-72 h. The aforementioned ionic liquid contains cations selected from N,N-dimethylimidazole ion, 1-ethyl-3-methylimidazole ion, 1-allyl-3-methylimidazole ion, 1-butyl-3-methylimidazole ion and 1-methyl-3-butylimidazole ion, and anions selected from chloride ion, bromide ion, acetate ion and thiocyanate ion. The method can be used to obtain enzymic hydrolysis rate of cellulose up to 100%.

L12 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:297680 CAPLUS

DOCUMENT NUMBER: 146:123861

TITLE: Dissolution of cellulose with ionic liquids and its application: a mini-review

AUTHOR(S): Zhu, Shengdong; Wu, Yuanxin; Chen, Qiming; Yu, Ziniu; Wang, Cunwen; Jin, Shiwei; Ding, Yigang; Wu, Gang

CORPORATE SOURCE: School of Chemical Engineering and Pharmacy, Hubei Key Laboratory of Novel Chemical Reactor and Green Chemical Technology, Wuhan Institute of Chemical Technology, Wuhan, 430073, Peop. Rep. China

SOURCE: Green Chemistry (2006), 8(4), 325-327

CODEN: GRCHFJ; ISSN: 1463-9262

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. In this paper, the dissoln. of cellulose with ionic liqs. and its application were reviewed.

Cellulose can be dissolved, without derivation, in some hydrophilic ionic liqs., such as 1-butyl-3-methylimidazolium chloride (BMIMCl) and 1-allyl-3-methylimidazolium chloride (AMIMCl). Microwave heating significantly accelerates the dissoln. process. Cellulose can be easily regenerated from its ionic liquid solns. by addition of water, ethanol or acetone. After its regeneration, the ionic liqs. can be recovered and reused. Fractionation of lignocellulosic materials and preparation of cellulose derivs. and composites are two of its typical applications. Although some basic studies, such as economical syntheses of ionic liqs. and studies of ionic liquid toxicol., are still much needed, commercialization of these processes has made great progress in recent years.

REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:523500 CAPLUS

DOCUMENT NUMBER: 143:28326

TITLE: Etherification of cellulose in ionic liquid solutions

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 23 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005054298	A1	20050616	WO 2004-FI730	20041202
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
FI 2003001763	A	20050604	FI 2003-1763	20031203
FI 116140	B1	20050930		
CA 2548007	A1	20050616	CA 2004-2548007	20041202
EP 1689788	A1	20060816	EP 2004-801227	20041202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS			
US 2007112185	A1	20070517	US 2007-581491	20070116
PRIORITY APPLN. INFO.:			FI 2003-1763	A 20031203
			WO 2004-FI730	W 20041202

OTHER SOURCE(S): MARPAT 143:28326

AB Cellulose is mixed and dissolved in an ionic liquid solvent and the solution is treated with an etherifying agent in the presence of inorg. base to form a cellulose ether, which is subsequently separated from the solution The dissoln. and the etherification are carried out in the absence of organic base and in the substantial absence of H₂O. Microwave irradiation and/or pressure can be applied to assist in dissoln. and etherification. Thus, 50 mg cellulose was dissolved in 5 g 1-butyl-3-methylimidazolium chloride (m. 60°) with the aid of microwaves to give 1% solution ClCH₂CO₂H (2.05 equiv) was added to the solution followed by 3.25 equiv of solid NaOH, the reaction mixture was heated for 2 h at 100° under microwave radiation and the resulting CM-cellulose was precipitated with MeOH, washed with MeOH and 80% aqueous MeOH, and dried.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:158715 CAPLUS

DOCUMENT NUMBER: 142:242565

TITLE: Dissolution and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005017001	A1	20050224	WO 2004-FI476	20040813
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2003001156	A	20050216	FI 2003-1156	20030815
FI 115835	B1	20050729		
CA 2532989	A1	20050224	CA 2004-2532989	20040813
EP 1654307	A1	20060510	EP 2004-742219	20040813
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004013435	A	20061010	BR 2004-13435	20040813
PRIORITY APPLN. INFO.:			FI 2003-1156	A 20030815
			WO 2004-FI476	W 20040813

OTHER SOURCE(S): MARPAT 142:242565

AB Wood, straw, and other natural lignocellulosic materials can be dissolved in an ionic liquid solvent under microwave irradiation and/or under pressure, and cellulose and other organic compds., such as lignin and extractives, can also be separated from the solution by precipitating with non-solvent , such as water, alcs., ketones, and ethers, of

cellulose. Thus, plywood sawdust was dissolved in 1-butyl-3-methyl-imidazolium chloride under microwave irradiation

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> "ionic liquid" and (carbohydrate or polysaccharide or sugar or cellulose or starch or chitin or pullulan or xanthan or dextran or glucan) adn microwave MISSING OPERATOR GLUCAN) ADN

The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> "ionic liquid" and (carbohydrate or polysaccharide or sugar or cellulose or starch or chitin or pullulan or xanthan or dextran or glucan) and microwave

284921 "IONIC"

511 "IONICS"

285185 "IONIC"

("IONIC" OR "IONICS")

800777 "LIQUID"

138530 "LIQUIDS"

904338 "LIQUID"

("LIQUID" OR "LIQUIDS")

1105042 "LIQ"

104871 "LIQS"

1145172 "LIQ"
 ("LIQ" OR "LIQS")
 1588658 "LIQUID"
 ("LIQUID" OR "LIQ")
 11079 "IONIC LIQUID"
 ("IONIC"(W) "LIQUID")
 133492 CARBOHYDRATE
 155004 CARBOHYDRATES
 224674 CARBOHYDRATE
 (CARBOHYDRATE OR CARBOHYDRATES)
 63148 POLYSACCHARIDE
 79822 POLYSACCHARIDES
 100435 POLYSACCHARIDE
 (POLYSACCHARIDE OR POLYSACCHARIDES)
 269177 SUGAR
 132316 SUGARS
 340895 SUGAR
 (SUGAR OR SUGARS)
 360710 CELLULOSE
 4428 CELLULOSES
 361213 CELLULOSE
 (CELLULOSE OR CELLULOSES)
 171072 STARCH
 9587 STARCHES
 172082 STARCH
 (STARCH OR STARCHES)
 17140 CHITIN
 313 CHITINS
 17155 CHITIN
 (CHITIN OR CHITINS)
 3654 PULLULAN
 1924 PULLULANS
 5169 PULLULAN
 (PULLULAN OR PULLULANS)
 13274 XANTHAN
 64 XANTHANS
 13278 XANTHAN
 (XANTHAN OR XANTHANS)
 37894 DEXTRAN
 4277 DEXTRANS
 38751 DEXTRAN
 (DEXTRAN OR DEXTRANS)
 15475 GLUCAN
 4421 GLUCANS
 16648 GLUCAN
 (GLUCAN OR GLUCANS)
 123811 MICROWAVE
 10771 MICROWAVES
 125649 MICROWAVE
 (MICROWAVE OR MICROWAVES)
 L13 9 "IONIC LIQUID" AND (CARBOHYDRATE OR POLYSACCHARIDE OR SUGAR OR
 CELLULOSE OR STARCH OR CHITIN OR PULLULAN OR XANTHAN OR DEXTRAN
 OR GLUCAN) AND MICROWAVE

=> d 113 1-9 ibib abs

L13 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:763873 CAPLUS
 DOCUMENT NUMBER: 147:141573
 TITLE: Processes for biomass treatment with higher energy efficiency
 INVENTOR(S): Gurin, Michael H.
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 21pp., Cont.-in-part of U.S. Ser. No. 309,025.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2007161095	A1	20070712	US 2007-691070	20070326
WO 2007112090	A2	20071004	WO 2007-US7430	20070326

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

PRIORITY APPLN. INFO.:
 US 2005-593485P P 20050118
 US 2005-595167P P 20050613
 US 2006-306911 A2 20060116
 US 2006-767403P P 20060325
 US 2006-309025 A2 20060612

AB A high efficiency method for synthesizing biomass fuels leveraging the synergistic impact of ionic liqs. on both the significant gains in pretreatment of biomass and the utilization of the combination of ionic liqs. and carbon dioxide under supercrit. conditions for energy generation is provided. The strategic use of heat exchangers, preferably microchannel heat exchangers and microchannel reactors further increase the efficiency and performance of the system by extensive heat recovery and the direct utilization of the biomass solution as the working fluid of a thermodyn. cycle.

L13 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:38602 CAPLUS
 DOCUMENT NUMBER: 146:123422
 TITLE: Ionic liquid reconstituted cellulose composites as solid support matrices with good transparency for biocatalytic reaction
 INVENTOR(S): Rogers, Robin D.; Daly, Daniel T.; Turner, Megan B.; Spear, Scott K.; Holbrey, John D.
 PATENT ASSIGNEE(S): The University of Alabama, USA
 SOURCE: PCT Int. Appl., 73pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent

LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2007005388	A2	20070111	WO 2006-US24863	20060627
WO 2007005388	A3	20070329		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CM, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

US 2007006774	A1	20070111	US 2006-475630	20060627
---------------	----	----------	----------------	----------

PRIORITY APPLN. INFO.: US 2005-694902P P 20050629

OTHER SOURCE(S): MARPAT 146:123422

AB Disclosed are composites comprising regenerated cellulose, a first active substance, a second active substance, and a linker. Thus, microcryst. cellulose was dissolved in 1-butyl-3-methylimidazolium chloride using microwave pulse heating at 120-150°, cooled to 60° to form a super-cooled liquid, 20% (based on cellulose) poly(L-lysine hydrobromide) was added therein, homogenized, cast onto a glass plate, the resulting film soaked in water for at least 24 h to leach residual from the film to give a reconstituted cellulose film, showing good transparency.

L13 ANSWER 3. OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:754555 CAPLUS

DOCUMENT NUMBER: 145:194651

TITLE: Method for complete enzymatic hydrolysis of straw cellulose pretreated with steam and microwave

INVENTOR(S): Chen, Hongzhang; Liu, Liying

PATENT ASSIGNEE(S): Institute of Process Engineering, Chinese Academy of Sciences, Peop. Rep. China

SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 7 pp.
CODEN: CNXXEV

DOCUMENT TYPE: Patent

LANGUAGE: Chinese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1806945	A	20060726	CN 2005-10011217	20050120
			CN 2005-10011217	20050120

PRIORITY APPLN. INFO.:

AB The title method comprises: (1) steam-blasting straws with water content of 10-35% under steam pressure of 1.0-1.5 MPa for 2-7 min, (2) washing with water of 50-100°C, drying, mixing with ionic liquid at a solid-liquid ratio of 1 : (5-50), and heating directly or

by microwave under stirring for 5-60 min, (3) washing the treated straw with water, and (4) hydrolyzing with cellulase at below 50°C and pH 4.8 for 48-72 h. The aforementioned ionic liquid contains cations selected from N,N-dimethylimidazole ion, 1-ethyl-3-methylimidazole ion, 1-allyl-3-methylimidazole ion, 1-butyl-3-methylimidazole ion and 1-methyl-3-butylimidazole ion, and anions selected from chloride ion, bromide ion, acetate ion and thiocyanate ion. The method can be used to obtain enzymic hydrolysis rate of cellulose up to 100%.

L13 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:559722 CAPLUS

DOCUMENT NUMBER: 146:338072

TITLE: Enzyme-catalyzed regioselective synthesis of sugar esters and related compounds

AUTHOR(S): Kennedy, John F.; Kumar, Harish; Panesar, Parmjit S.; Marwaha, Satwinder S.; Goyal, Rita; Parmar, Anupama; Kaur, Sukhwinder

CORPORATE SOURCE: Birmingham Carbohydrate and Protein Technology Group, School of Chemistry, University of Birmingham, Birmingham, B15 2TT, UK

SOURCE: Journal of Chemical Technology and Biotechnology (2006), 81(6), 866-876
CODEN: JCTBED; ISSN: 0268-2575

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB In this review, a comprehensive and illustrative survey is made of the regioselective synthesis of esters of sugars and related compds. using lipases. The main emphasis has been given to the screening and use of com. available lipases for the enzymic esterification of neutral monosaccharides, disaccharides, sugar alcs. and their selected ether and ester derivs. The effect of solvents and solubilizing agents in improving the yields of the resultant sugar fatty acid esters has been incorporated. Further, solvent-free esterification with molten fatty acids, use of ionic liqs. and microwave radiations for improvement in the methodol. have also been discussed.

REFERENCE COUNT: 79 THERE ARE 79 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:297680 CAPLUS

DOCUMENT NUMBER: 146:123861

TITLE: Dissolution of cellulose with ionic liquids and its application: a mini-review

AUTHOR(S): Zhu, Shengdong; Wu, Yuanxin; Chen, Qiming; Yu, Ziniu; Wang, Cunwen; Jin, Shiwei; Ding, Yigang; Wu, Gang

CORPORATE SOURCE: School of Chemical Engineering and Pharmacy, Hubei Key Laboratory of Novel Chemical Reactor and Green Chemical Technology, Wuhan Institute of Chemical Technology, Wuhan, 430073, Peop. Rep. China

SOURCE: Green Chemistry (2006), 8(4), 325-327

CODEN: GRCHFJ; ISSN: 1463-9262

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. In this paper, the dissoln. of cellulose with

Cellulose can be dissolved, without derivation, in some hydrophilic ionic liqs., such as 1-butyl-3-methylimidazolium chloride (BMIMCl) and 1-allyl-3-methylimidazolium chloride (AMIMCl). Microwave heating significantly accelerates the dissoln. process. Cellulose can be easily regenerated from its ionic liquid solns. by addition of water, ethanol or acetone. After its regeneration, the ionic liqs. can be recovered and reused. Fractionation of lignocellulosic materials and preparation of cellulose derivs. and composites are two of its typical applications. Although some basic studies, such as economical syntheses of ionic liqs. and studies of ionic liquid toxicol., are still much needed, commercialization of these processes has made great progress in recent years.

L13 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

PATENT INFORMATION:

AB Cellulose is mixed and dissolved in an ionic liquid solvent and the solution is treated with an etherifying agent in

the presence of inorg. base to form a cellulose ether, which is subsequently separated from the solution The dissoln. and the etherification are

carried out in the absence of organic base and in the substantial absence of H₂O. Microwave irradiation and/or pressure can be applied to assist in dissoln. and etherification. Thus, 50 mg cellulose was dissolved in 5 g 1-butyl-3-methylimidazolium chloride (m. 60°) with the aid of microwaves to give 1% solution ClCH₂CO₂H (2.05 equiv) was added to the solution followed by 3.25 equiv of solid NaOH, the reaction mixture was heated for 2 h at 100° under microwave radiation and the resulting CM-cellulose was precipitated with MeOH, washed with MeOH and 80% aqueous MeOH, and dried.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:239036 CAPLUS

DOCUMENT NUMBER: 142:299721

TITLE: Esterification of starch under microwave irradiation and pressure

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005023873	A1	20050317	WO 2004-FI523	20040910
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
FI 2003001301	A	20050312	FI 2003-1301	20030911
FI 116142	B1	20050930		
CA 2533553	A1	20050317	CA 2004-2533553	20040910
EP 1664125	A1	20060607	EP 2004-767037	20040910
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
BR 2004013432	A	20061010	BR 2004-13432	20040910
US 2007073051	A1	20070329	US 2006-566975	20061207
PRIORITY APPLN. INFO.:			FI 2003-1301	A 20030911
			WO 2004-FI523	W 20040910

OTHER SOURCE(S): MARPAT 142:299721

AB An organic starch ester is prepared by mixing a starch material, such as natural starch or hydrolyzed starch, with an ionic liquid solvent to dissolve the

starch, and then treating the dissolved starch with an organic esterifying agent, such as C1-11 carboxylic acid, to form an organic starch ester, and subsequently separating the organic starch ester from the solution by adding a non-solvent, such as alcs., ketones, and acetonitrile, to the starch ester solution Microwave irradiation and/or pressure can be applied to assist the dissoln. and esterification. Thus, native barely starch was dissolved in ionic 1-butyl-3-methylimidazolium chloride and reacted with acetic anhydride, followed by quenching with ethanol to receive starch acetate.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:158715 CAPLUS

DOCUMENT NUMBER: 142:242565

TITLE: Dissolution and delignification of lignocellulosic materials with ionic liquid solvent under microwave irradiation

INVENTOR(S): Myllymaeki, Vesa; Aksela, Reijo

PATENT ASSIGNEE(S): Kemira Oyj, Finland

SOURCE: PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005017001	A1	20050224	WO 2004-FI476	20040813
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2003001156	A	20050216	FI 2003-1156	20030815
FI 115835	B1	20050729		
CA 2532989	A1	20050224	CA 2004-2532989	20040813
EP 1654307	A1	20060510	EP 2004-742219	20040813
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004013435	A	20061010	BR 2004-13435	20040813
PRIORITY APPLN. INFO.:			FI 2003-1156	A 20030815
			WO 2004-FI476	W 20040813

OTHER SOURCE(S): MARPAT 142:242565

AB Wood, straw, and other natural lignocellulosic materials can be dissolved in an ionic liquid solvent under microwave irradiation and/or under pressure, and cellulose and other organic compds., such as lignin and extractives, can also be separated from the solution

by precipitating with non-solvent , such as water, alcs., ketones, and ethers,
of

cellulose. Thus, plywood sawdust was dissolved in
1-butyl-3-methyl-imidazolium chloride under microwave irradiation

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:653978 CAPLUS

DOCUMENT NUMBER: 132:219083

TITLE: Attempted dissolution of "insoluble organic matter"
from Methanococcus jannaschii and Rostherne Mere (UK)
sediment with 1-ethyl-3-methyl imidazolium
chloride/aluminum (III) chloride

AUTHOR(S): Sutton, P. A.; Lewis, C. A.; Patell, Y.; Seddon, K.
R.; Rowland, S. J.

CORPORATE SOURCE: Petroleum and Environmental Geochemistry Group,
Department of Environmental Sciences, University of
Plymouth, Plymouth, PL4 8AA, UK

SOURCE: Ancient Biomolecules (1998), 2(2-3), 195-207
CODEN: ANBIFP; ISSN: 1358-6122

PUBLISHER: Harwood Academic Publishers

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Despite its geochem. significance, sedimentary macromol. organic matter,
including biomol. material, which is insol. in non-oxidizing acids, bases
and normal organic solvents (insol. organic matter, IOM), and which is termed
kerogen when it occurs in Ancient sediments, has only been characterized
in a limited number of elegant chemolytic studies. This owes largely to its
insoly., which reduces the number of anal. methods applicable for structural
elucidation. In the present, preliminary study we report our attempts to
dissolve IOM in order to make it more amenable to anal. by conventional
methods. IOM was isolated by sequential removal of soluble matter from
Methanococcus jannaschii (2.8% \pm 0.3% dry weight IOM, n = 3) and from a
Recent lacustrine sediment (Rostherne Mere, UK, 11-15% dry weight IOM). For
the sediment, the sequential removal of proteins, carbohydrates,
lipids and minerals was monitored by examination of the insol. residues by
pyrolysis-gas chromatog./mass spectrometry (py-GC/MS) and solid state NMR
spectroscopy (ss-NMR). IOM samples were then treated with an
ionic liquid, 1-ethyl-3-Me imidazolium chloride/aluminum
(III) chloride ([emim]Cl/AlCl₃). A synthetic dendrimer was also treated
with ionic liquid in order to assess any effects on a
fairly high mol. weight (1075), but non-macromol., substrate of known
structure which contains chemical moieties which have been identified in
kerogens by chemolytic studies (e.g. aromatic rings, ether and ester
linkages, and methylene groups). Ionic liquid/sample
mixts. were microwave heated, hydrolyzed, filtered and both
non-aqueous soluble retentates and aqueous filtrates collected. Retentates
and aqueous

filtrates were then extracted with organic solvents. The insol. matter in the
retentates was examined by py-GC/MS. The organic solvent exts. of the
retentates and of the aqueous filtrates were investigated using solution NMR,
much of which remains to be identified by interpretation of the
spectroscopic data. However, some preliminary observations can be made.
Interestingly, the proton NMR spectrum of the dendrimer filtrate extract
showed that ionic liquid treatment had substantially
changed the dendrimer structure. Over 30 new alkyl resonances appeared in

the aliphatic region of the spectrum (δ 0.5-4.0 ppm), and the pattern of aromatic resonances (δ 6.3-8.2 ppm) was also changed. It is suggested that Friedel-Crafts-type alkylations could have been promoted by the aluminum (III) chloride moiety of the [emim]Cl/AlCl₃. When applied to unknown IOM from bacteria and sediments it appears that the addition of ionic liquid, heating and hydrolysis also chemical alters these substrates, since some previously solvent-insol. organic matter became soluble. The proton NMR spectrum of the IOM filtrate extract of *M. jannaschii* contained predominantly aliphatic resonances with only one aromatic and two alkenic resonances. Proton NMR spectra of the sedimentary IOM filtrate exts. were also dominated by aliphatic signals, in contrast to solid state NMR anal. of the IOM before treatment, again suggesting alkylation of the original matter. If the mode of action of ionic liquid treatment is reproducible and can be better understood, this procedure may play a part in the further characterization of IOM, but simple dissoln. of IOM appears not to occur, judging from these preliminary expts.

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> fiel stng

30 FIEL
3 FIELDS
33 FIEL
(FIEL OR FIELDS)
2 STNG
L14 0 FIEL STNG
(FIEL(W) STNG)

=> file stng

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	114.92	235.20
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-17.94	-42.12

FILE 'STNGUIDE' ENTERED AT 16:06:09 ON 26 NOV 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Nov 23, 2007 (20071123/UP).

=> d his

(FILE 'HOME' ENTERED AT 15:58:30 ON 26 NOV 2007)

FILE 'CAPLUS' ENTERED AT 15:58:43 ON 26 NOV 2007

E MYLLYMAKI/AU

L1 19 S E2
E AKSELA/AU
L2 47 S E10 OR E11
L3 60 DUP REMOVE L1 L2 (6 DUPLICATES REMOVED)
L4 19 S L3
L5 41 S L3

L6

6 S L3 AND IONIC

FILE 'STNGUIDE' ENTERED AT 16:00:24 ON 26 NOV 2007

FILE 'CAPLUS' ENTERED AT 16:00:34 ON 26 NOV 2007

E SWATLOSKI/AU

L7 91 S E4-E7

L8 83 S L7 AND "IONIC LIQUID"

L9 26 S L8 AND CELLULOSE

FILE 'STNGUIDE' ENTERED AT 16:01:40 ON 26 NOV 2007

FILE 'CAPLUS' ENTERED AT 16:03:16 ON 26 NOV 2007

L10 177 "IONIC LIQUID" AND CELLULOSE

L11 11 L10 AND ETHER?

L12 6 L10 AND MICROWAVE

L13 9 "IONIC LIQUID" AND (CARBOHYDRATE OR POLYSACCHARIDE OR SUGAR OR

L14 0 FIEL STNG

FILE 'STNGUIDE' ENTERED AT 16:06:09 ON 26 NOV 2007

=>

---Logging off of STN---

=